

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/001420

International filing date: 12 February 2005 (12.02.2005)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 04004326.7
Filing date: 26 February 2004 (26.02.2004)

Date of receipt at the International Bureau: 13 April 2005 (13.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

EP/05/1420

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04004326.7

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 04004326.7
Demande no:

Anmeldetag:
Date of filing: 26.02.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Bayer HealthCare AG
Kaiser-Wilhelm-Allee
51373 Leverkusen
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Diagnostics and therapeutics for diseases associated with glycogen synthase
kinase 3 beta (GSK3B)

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

G01N33/48

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

26. Feb. 2004

Diagnostics and Therapeutics for Diseases Associated with Glycogen Synthase Kinase 3 beta (GSK3B)

Technical field of the invention

5 The present invention is in the field of molecular biology, more particularly, the present invention relates to nucleic acid sequences and amino acid sequences of a human GSK3B and its regulation for the treatment of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in mammals.

Background of the invention

10 GSK3B is a member of the enzyme group of kinases [[Woodgett (1990)], [Stambolic and Woodgett, (1994)], [Plyte et al., (1992)], [Frame et al. (2001)], US 6500938, US 6323029, WO 03068961]. Kinases are enzymes, which catalyse the transfer of a phosphate group (phosphorylation) from a donor (mainly ATP) onto an acceptor molecule's nucleophilic functional group, such as hydroxy-, carboxy-, guanidino-, thiol-, or imidazole- groups. Kinases can act on on a variety of molecules like small metabolites, or proteins (protein kinases).

15 Kinases that phosphorylate small organic molecules often play important roles in metabolic pathways such as glycolysis (e.g. hexokinase, glucokinase, or phosphofructokinase), or in anabolic pathways.

Protein kinases play critical roles in cellular signal transduction.

20 Cellular signal transduction is a fundamental mechanism whereby extracellular stimuli are relayed to the interior of cells and subsequently regulate diverse cellular processes. One of the key biochemical mechanisms of signal transduction involves the reversible phosphorylation of proteins. Phosphorylation of polypeptides regulates the activity of mature proteins by altering their structure and function. Phosphate most often resides on the hydroxyl moiety (—OH) of serine, threonine, or tyrosine amino acids in proteins.

25 Kinases regulate many different cell proliferation, differentiation, and signaling processes by adding phosphate groups to proteins. Uncontrolled signaling has been implicated in a variety of disease conditions including inflammation, cancer, arteriosclerosis, and psoriasis. Reversible protein phosphorylation is the main strategy for controlling activities of eukaryotic cells. The high energy phosphate, which drives activation, is generally transferred from adenosine triphosphate
30 molecules (ATP) to a particular protein by protein kinases and removed from that protein by protein phosphatases. Phosphorylation occurs in response to extracellular signals (hormones,

neurotransmitters, growth and differentiation factors, etc), cell cycle checkpoints, and environmental or nutritional stresses and is roughly analogous to turning on a molecular switch. When the switch goes on, the appropriate protein kinase activates a metabolic enzyme, regulatory protein, receptor, cytoskeletal protein, ion channel or pump, or transcription factor.

- 5 The kinases comprise the largest known protein group, a superfamily of enzymes with widely varied functions and specificities. They are usually named after their substrate, their regulatory molecules, or some aspect of a mutant phenotype. With regard to substrates, the protein kinases may be roughly divided into two groups; those that phosphorylate tyrosine residues (protein tyrosine kinases, PTK) and those that phosphorylate serine or threonine residues (serine/threonine
- 10 kinases, STK). A few protein kinases have dual specificity and phosphorylate threonine and tyrosine residues. Almost all kinases contain a similar 250-300 amino acid catalytic domain. The N-terminal domain, which contains subdomains I-IV, generally folds into a two-lobed structure, which binds and orients the ATP (or GTP) donor molecule. The larger C terminal lobe, which contains subdomains VI-XI, binds the protein substrate and carries out the transfer of the gamma
- 15 phosphate from ATP to the hydroxyl group of a serine, threonine, or tyrosine residue. Subdomain V spans the two lobes.

The kinases may be categorized into families by the different amino acid sequences (generally between 5 and 100 residues) located on either side of, or inserted into loops of, the kinase domain. These added amino acid sequences allow the regulation of each kinase as it recognizes and

20 interacts with its target protein. The primary structure of the kinase domains is conserved and can be further subdivided into 11 subdomains. Each of the 11 subdomains contains specific residues and motifs or patterns of amino acids that are characteristic of that subdomain and are highly conserved [Hardie, G. and Hanks, S. (1995)].

The second messenger dependent protein kinases primarily mediate the effects of second

25 messengers such as cyclic AMP (cAMP), cyclic GMP, inositol triphosphate, phosphatidylinositol, 3,4,5-triphosphate, cyclic-ADPribose, arachidonic acid, diacylglycerol and calcium-calmodulin. The cyclic-AMP dependent protein kinases (PKA) are important members of the STK family. Cyclic-AMP is an intracellular mediator of hormone action in all prokaryotic and animal cells that have been studied. Such hormone-induced cellular responses include thyroid hormone secretion,

30 cortisol secretion, progesterone secretion, glycogen breakdown, bone resorption, and regulation of heart rate and force of heart muscle contraction. PKA is found in all animal cells and is thought to account for the effects of cyclic-AMP in most of these cells. Altered PKA expression is implicated in a variety of disorders and diseases including cancer, thyroid disorders, diabetes, atherosclerosis, and cardiovascular disease [Isselbacher, K. J. et al. (1994)].

Calcium-calmodulin (CaM) dependent protein kinases are also members of STK family. Calmodulin is a calcium receptor that mediates many calcium regulated processes by binding to target proteins in response to the binding of calcium. The principle target protein in these processes is CaM dependent protein kinases. CaM-kinases are involved in regulation of smooth muscle contraction (MLC kinase), glycogen breakdown (phosphorylase kinase); and neuro-transmission (CaM kinase I and CaM kinase II). CaM kinase I phosphorylates a variety of substrates including the neurotransmitter related proteins synapsin I and II, the gene transcription regulator, CREB, and the cystic fibrosis conductance regulator protein, CFTR [Haribabu, B. et al. (1995)]. CaM II kinase also phosphorylates synapsin at different sites, and controls the synthesis of catecholamines in the brain through phosphorylation and activation of tyrosine hydroxylase. Many of the CaM kinases are activated by phosphorylation in addition to binding to CaM. The kinase may autophosphorylate itself, or be phosphorylated by another kinase as part of a "kinase cascade".

Another ligand-activated protein kinase is 5'-AMP-activated protein kinase (AMPK) [Gao, G. et al. (1996)]. Mammalian AMPK is a regulator of fatty acid and sterol synthesis through phosphorylation of the enzymes acetyl-CoA carboxylase and hydroxymethylglutaryl-CoA reductase and mediates responses of these pathways to cellular stresses such as heat shock and depletion of glucose and ATP. AMPK is a heterotrimeric complex comprised of a catalytic alpha subunit and two non-catalytic beta and gamma subunits that are believed to regulate the activity of the alpha subunit. Subunits of AMPK have a much wider distribution in non-lipogenic tissues such as brain, heart, spleen, and lung than expected. This distribution suggests that its role may extend beyond regulation of lipid metabolism alone.

The mitogen-activated protein kinases (MAP) are also members of the STK family. MAP kinases also regulate intracellular signaling pathways. They mediate signal transduction from the cell surface to the nucleus via phosphorylation cascades. Several subgroups have been identified, and each manifests different substrate specificities and responds to distinct extracellular stimuli [Egan, S. E. and Weinberg, R. A. (1993)]. MAP kinase signaling pathways are present in mammalian cells as well as in yeast. The extracellular stimuli that activate mammalian pathways include epidermal growth factor (EGF), ultraviolet light, hyperosmolar medium, heat shock, endotoxic lipopolysaccharide (LPS), and pro-inflammatory cytokines such as tumor necrosis factor (TNF) and interleukin-1 (IL-1).

PRK (proliferation-related kinase) is a serum/cytokine inducible STK that is involved in regulation of the cell cycle and cell proliferation in human megakaryotic cells [Li, B. et al. (1996)]. PRK is related to the polo (derived from humans polo gene) family of STKs implicated in cell division.

PRK is downregulated in lung tumor tissue and may be a proto-oncogene whose deregulated expression in normal tissue leads to oncogenic transformation. Altered MAP kinase expression is implicated in a variety of disease conditions including cancer, inflammation, immune disorders, and disorders affecting growth and development.

- 5 The cyclin-dependent protein kinases (CDKs) are another group of STKs that control the progression of cells through the cell cycle. Cyclins are small regulatory proteins that act by binding to and activating CDKs that then trigger various phases of the cell cycle by phosphorylating and activating selected proteins involved in the mitotic process. CDKs are unique in that they require multiple inputs to become activated. In addition to the binding of cyclin, CDK activation requires
- 10 the phosphorylation of a specific threonine residue and the dephosphorylation of a specific tyrosine residue.

- Protein tyrosine kinases, PTKs, specifically phosphorylate tyrosine residues on their target proteins and may be divided into transmembrane, receptor PTKs and nontransmembrane, non-receptor PTKs. Transmembrane protein-tyrosine kinases are receptors for most growth factors.
- 15 Binding of growth factor to the receptor activates the transfer of a phosphate group from ATP to selected tyrosine side chains of the receptor and other specific proteins. Growth factors (GF) associated with receptor PTKs include; epidermal GF, platelet-derived GF, fibroblast GF, hepatocyte GF, insulin and insulin-like GFs, nerve GF, vascular endothelial GF, and macrophage colony stimulating factor.
- 20 Non-receptor PTKs lack transmembrane regions and, instead, form complexes with the intracellular regions of cell surface receptors. Such receptors that function through non-receptor PTKs include those for cytokines, hormones (growth hormone and prolactin) and antigen-specific receptors on T and B lymphocytes.

- In an effort to discover novel treatments for diseases, biomedical researchers and chemists have designed, synthesized, and tested molecules that inhibit the function of protein kinases. Some
- 25 small organic molecules form a class of compounds that modulate the function of protein kinases. The modulatory compounds are potentially advantageous therapeutics for disease conditions including but not limited to inflammation, cancer, arteriosclerosis, and psoriasis.

TaqMan-Technology / expression profiling

- 30 TaqMan is a recently developed technique, in which the release of a fluorescent reporter dye from a hybridisation probe in real-time during a polymerase chain reaction (PCR) is proportional to the accumulation of the PCR product. Quantification is based on the early, linear part of the reaction,

and by determining the threshold cycle (CT), at which fluorescence above background is first detected.

Gene expression technologies may be useful in several areas of drug discovery and development, such as target identification, lead optimization, and identification of mechanisms of action. The TaqMan technology can be used to compare differences between expression profiles of normal tissue and diseased tissue. Expression profiling has been used in identifying genes, which are up- or downregulated in a variety of diseases. An interesting application of expression profiling is temporal monitoring of changes in gene expression during disease progression and drug treatment or in patients versus healthy individuals. The premise in this approach is that changes in pattern of gene expression in response to physiological or environmental stimuli (e.g., drugs) may serve as indirect clues about disease-causing genes or drug targets. Moreover, the effects of drugs with established efficacy on global gene expression patterns may provide a guidepost, or a genetic signature, against which a new drug candidate can be compared.

GSK3B

The nucleotide sequence of GSK3B is accessible in the databases by the accession number BC000251 and is given in SEQ ID NO:1. The amino acid sequence of GSK3B depicted in SEQ ID NO:2.

Glycogen synthase kinase-3 (GSK3) is a proline-directed serine-threonine kinase that was initially identified as a phosphorylating and inactivating glycogen synthase. Two isoforms, alpha (GSK3A) and beta, show a high degree of amino acid homology [Stambolic and Woodgett, (1994)]. GSK3B is involved in energy metabolism, neuronal cell development, and body pattern formation [Plyte et al., (1992)].

Woodgett [Woodgett (1990)] cloned rat Gsk3a and GSK3B. The deduced 483-amino acid Gsk3a protein is 93% identical overall and 99% identical in the kinase catalytic domain to the human protein. SDS-PAGE analysis showed expression of the 51-kD rat protein as predicted from the primary sequence. Northern blot analysis revealed wide expression of a 2.5-kb transcript in rat tissues. Western blot analysis, however, showed that expression is variable, suggesting differential modes of transcriptional and translational regulation.

Frame et al. [Frame et al. (2001)] demonstrated that the insulin-induced inhibition of GSK3 and its unique substrate specificity are explained by the existence of a phosphate-binding site in which arg96 is critical. Mutation of arg96 abolished the phosphorylation of 'primed' glycogen synthase as well as inhibition by protein kinase B-mediated phosphorylation of ser9. Hence, the phosphory-

lated N terminus acts as a pseudosubstrate, occupying the same phosphate-binding site used by primed substrates. This mutation did not affect phosphorylation of 'nonprimed' substrates in the Wnt-signaling pathway (axin and beta-catenin), suggesting novel approaches to design more selective GSK3 inhibitors for the treatment of diabetes.

- 5 GSK3B is published in patents US 6500938, US 6323029 and WO03068961.

Summary of the invention

The invention relates to novel disease associations of GSK3B polypeptides and polynucleotides. The invention also relates to novel methods of screening for therapeutic agents for the treatment of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation,
10 respiratory diseases, neurological diseases and urological diseases in a mammal. The invention also relates to pharmaceutical compositions for the treatment of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a GSK3B polypeptide, a GSK3B polynucleotide, or regulators of GSK3B or modulators of GSK3B activity. The invention further
15 comprises methods of diagnosing cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal.

Brief Description of the Drawings

Fig. 1 shows the nucleotide sequence of a GSK3B polynucleotide (SEQ ID NO:1).

- 20 Fig. 2 shows the amino acid sequence of a GSK3B polypeptide (SEQ ID NO:2).

Fig. 3 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO:3).

Fig. 4 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO:4).

Fig. 5 shows a nucleotide sequence useful as a probe to detect proteins of the invention (SEQ ID NO:5).

25 Detailed description of the invention

Definition of terms

An "oligonucleotide" is a stretch of nucleotide residues which has a sufficient number of bases to be used as an oligomer, amplimer or probe in a polymerase chain reaction (PCR). Oligonucleotides are prepared from genomic or cDNA sequence and are used to amplify, reveal, or

confirm the presence of a similar DNA or RNA in a particular cell or tissue. Oligonucleotides or oligomers comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 35 nucleotides, preferably about 25 nucleotides.

5 "Probes" may be derived from naturally occurring or recombinant single- or double-stranded nucleic acids or may be chemically synthesized. They are useful in detecting the presence of identical or similar sequences. Such probes may be labeled with reporter molecules using nick translation, Klenow fill-in reaction, PCR or other methods well known in the art. Nucleic acid probes may be used in southern, northern or in situ hybridizations to determine whether DNA or RNA encoding a certain protein is present in a cell type, tissue, or organ.

10 A "fragment of a polynucleotide" is a nucleic acid that comprises all or any part of a given nucleotide molecule, the fragment having fewer nucleotides than about 6 kb, preferably fewer than about 1 kb.

15 "Reporter molecules" are radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents which associate with a particular nucleotide or amino acid sequence, thereby establishing the presence of a certain sequence, or allowing for the quantification of a certain sequence.

"Chimeric" molecules may be constructed by introducing all or part of the nucleotide sequence of this invention into a vector containing additional nucleic acid sequence which might be expected to change any one or several of the following GSK3B characteristics: cellular location, distribution, ligand-binding affinities, interchain affinities, degradation/turnover rate, signaling, etc.

20 "Active", with respect to a GSK3B polypeptide, refers to those forms, fragments, or domains of a GSK3B polypeptide which retain the biological and/or antigenic activity of a GSK3B polypeptide.

25 "Naturally occurring GSK3B polypeptide" refers to a polypeptide produced by cells which have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including but not limited to acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

"Derivative" refers to polypeptides which have been chemically modified by techniques such as ubiquitination, labeling (see above), pegylation (derivatization with polyethylene glycol), and chemical insertion or substitution of amino acids such as ornithine which do not normally occur in human proteins.

"Conservative amino acid substitutions" result from replacing one amino acid with another having similar structural and/or chemical properties, such as the replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

5 "Insertions" or "deletions" are typically in the range of about 1 to 5 amino acids. The variation allowed may be experimentally determined by producing the peptide synthetically while systematically making insertions, deletions, or substitutions of nucleotides in the sequence using recombinant DNA techniques.

10 A "signal sequence" or "leader sequence" can be used, when desired, to direct the polypeptide through a membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous sources by recombinant DNA techniques.

An "oligopeptide" is a short stretch of amino acid residues and may be expressed from an oligonucleotide. Oligopeptides comprise a stretch of amino acid residues of at least 3, 5, 10 amino acids and at most 10, 15, 25 amino acids, typically of at least 9 to 13 amino acids, and of sufficient length to display biological and/or antigenic activity.

15 "Inhibitor" is any substance which retards or prevents a chemical or physiological reaction or response. Common inhibitors include but are not limited to antisense molecules, antibodies, and antagonists.

20 "Standard expression" is a quantitative or qualitative measurement for comparison. It is based on a statistically appropriate number of normal samples and is created to use as a basis of comparison when performing diagnostic assays, running clinical trials, or following patient treatment profiles.

"Animal" as used herein may be defined to include human, domestic (e.g., cats, dogs, etc.), agricultural (e.g., cows, horses, sheep, etc.) or test species (e.g., mouse, rat, rabbit, etc.).

A "GSK3B polynucleotide", within the meaning of the invention, shall be understood as being a nucleic acid molecule selected from a group consisting of

- 25 (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2,
- (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1,
- (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1,

- (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and
 - (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code;
- 5 wherein the polypeptide encoded by said nucleic acid molecule has GSK3B activity.

A "GSK3B polypeptide", within the meaning of the invention, shall be understood as being a polypeptide selected from a group consisting of

- (i) polypeptides having the sequence of SEQ ID NO: 2,
- (ii) polypeptides comprising the sequence of SEQ ID NO: 2,
- 10 (iii) polypeptides encoded by GSK3B polynucleotides; and
- (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii);

wherein said polypeptide has GSK3B activity.

- 15 The nucleotide sequences encoding a GSK3B (or their complement) have numerous applications in techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use in the construction of oligomers for PCR, use for chromosome and gene mapping, use in the recombinant production of GSK3B, and use in generation of antisense DNA or RNA, their chemical analogs and the like. Uses of nucleotides encoding a GSK3B disclosed herein are exemplary of known techniques and are not intended to limit their use in any
- 20 technique known to a person of ordinary skill in the art. Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, e.g., the triplet genetic code, specific base pair interactions, etc.

- 25 It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of GSK3B - encoding nucleotide sequences may be produced. Some of these will only bear minimal homology to the nucleotide sequence of the known and naturally occurring GSK3B. The invention has specifically contemplated each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as

applied to the nucleotide sequence of naturally occurring GSK3B, and all such variations are to be considered as being specifically disclosed.

Although the nucleotide sequences which encode a GSK3B, its derivatives or its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring GSK3B polynucleotide under stringent conditions, it may be advantageous to produce nucleotide sequences encoding GSK3B polypeptides or its derivatives possessing a substantially different codon usage. Codons can be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding a GSK3B polypeptide and/or its derivatives without altering the encoded amino acid sequence include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

Nucleotide sequences encoding a GSK3B polypeptide may be joined to a variety of other nucleotide sequences by means of well established recombinant DNA techniques. Useful nucleotide sequences for joining to GSK3B polynucleotides include an assortment of cloning vectors such as plasmids, cosmids, lambda phage derivatives, phagemids, and the like. Vectors of interest include expression vectors, replication vectors, probe generation vectors, sequencing vectors, etc. In general, vectors of interest may contain an origin of replication functional in at least one organism, convenient restriction endonuclease sensitive sites, and selectable markers for one or more host cell systems.

Another aspect of the subject invention is to provide for GSK3B-specific hybridization probes capable of hybridizing with naturally occurring nucleotide sequences encoding GSK3B. Such probes may also be used for the detection of similar kinase encoding sequences and should preferably show at least 40% nucleotide identity to GSK3B polynucleotides. The hybridization probes of the subject invention may be derived from the nucleotide sequence presented as SEQ ID NO: 1 or from genomic sequences including promoter, enhancers or introns of the native gene. Hybridization probes may be labelled by a variety of reporter molecules using techniques well known in the art.

It will be recognized that many deletional or mutational analogs of GSK3B polynucleotides will be effective hybridization probes for GSK3B polynucleotides. Accordingly, the invention relates to nucleic acid sequences that hybridize with such GSK3B encoding nucleic acid sequences under stringent conditions.

"Stringent conditions" refers to conditions that allow for the hybridization of substantially related nucleic acid sequences. For instance, such conditions will generally allow hybridization of sequence with at least about 85% sequence identity, preferably with at least about 90% sequence identity, more preferably with at least about 95% sequence identity. Hybridization conditions and probes can be adjusted in well-characterized ways to achieve selective hybridization of human-derived probes. Stringent conditions, within the meaning of the invention are 65°C in a buffer containing 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7 % (w/v) SDS.

Nucleic acid molecules that will hybridize to GSK3B polynucleotides under stringent conditions can be identified functionally. Without limitation, examples of the uses for hybridization probes include: histochemical uses such as identifying tissues that express GSK3B; measuring mRNA levels, for instance to identify a sample's tissue type or to identify cells that express abnormal levels of GSK3B; and detecting polymorphisms of GSK3B.

PCR provides additional uses for oligonucleotides based upon the nucleotide sequence which encodes GSK3B. Such probes used in PCR may be of recombinant origin, chemically synthesized, or a mixture of both. Oligomers may comprise discrete nucleotide sequences employed under optimized conditions for identification of GSK3B in specific tissues or diagnostic use. The same two oligomers, a nested set of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for identification of closely related DNAs or RNAs.

Rules for designing polymerase chain reaction (PCR) primers are now established, as reviewed by PCR Protocols. Degenerate primers, i.e., preparations of primers that are heterogeneous at given sequence locations, can be designed to amplify nucleic acid sequences that are highly homologous to, but not identical with GSK3B. Strategies are now available that allow for only one of the primers to be required to specifically hybridize with a known sequence. For example, appropriate nucleic acid primers can be ligated to the nucleic acid sought to be amplified to provide the hybridization partner for one of the primers. In this way, only one of the primers need be based on the sequence of the nucleic acid sought to be amplified.

PCR methods for amplifying nucleic acid will utilize at least two primers. One of these primers will be capable of hybridizing to a first strand of the nucleic acid to be amplified and of priming enzyme-driven nucleic acid synthesis in a first direction. The other will be capable of hybridizing the reciprocal sequence of the first strand (if the sequence to be amplified is single stranded, this sequence will initially be hypothetical, but will be synthesized in the first amplification cycle) and of priming nucleic acid synthesis from that strand in the direction opposite the first direction and towards the site of hybridization for the first primer. Conditions for conducting such amplifications, particularly under preferred stringent hybridization conditions, are well known.

Other means of producing specific hybridization probes for GSK3B include the cloning of nucleic acid sequences encoding GSK3B or GSK3B derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate reporter molecules.

It is possible to produce a DNA sequence, or portions thereof, entirely by synthetic chemistry. After synthesis, the nucleic acid sequence can be inserted into any of the many available DNA vectors and their respective host cells using techniques which are well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into the nucleotide sequence. Alternately, a portion of sequence in which a mutation is desired can be synthesized and recombined with longer portion of an existing genomic or recombinant sequence.

GSK3B polynucleotides may be used to produce a purified oligo-or polypeptide using well known methods of recombinant DNA technology. The oligopeptide may be expressed in a variety of host cells, either prokaryotic or eukaryotic. Host cells may be from the same species from which the nucleotide sequence was derived or from a different species. Advantages of producing an oligonucleotide by recombinant DNA technology include obtaining adequate amounts of the protein for purification and the availability of simplified purification procedures.

Quantitative determinations of nucleic acids

An important step in the molecular genetic analysis of human disease is often the enumeration of the copy number of a nucleic acid or the relative expression of a gene in particular tissues.

Several different approaches are currently available to make quantitative determinations of nucleic acids. Chromosome-based techniques, such as comparative genomic hybridization (CGH) and fluorescent in situ hybridization (FISH) facilitate efforts to cytogenetically localize genomic regions that are altered in tumor cells. Regions of genomic alteration can be narrowed further using loss of heterozygosity analysis (LOH), in which disease DNA is analyzed and compared with normal DNA for the loss of a heterozygous polymorphic marker. The first experiments used restriction fragment length polymorphisms (RFLPs) [Johnson, (1989)], or hypervariable minisatellite DNA [Barnes, 2000]. In recent years LOH has been performed primarily using PCR amplification of microsatellite markers and electrophoresis of the radio labelled [Jeffreys, (1985)] or fluorescently labelled PCR products [Weber, (1990)] and compared between paired normal and disease DNAs.

A number of other methods have also been developed to quantify nucleic acids [Gergen, (1992)]. More recently, PCR and RT-PCR methods have been developed which are capable of measuring the amount of a nucleic acid in a sample. One approach, for example, measures PCR product quantity in the log phase of the reaction before the formation of reaction products plateaus
5 [Thomas, (1980)].

A gene sequence contained in all samples at relatively constant quantity is typically utilized for sample amplification efficiency normalization. This approach, however, suffers from several drawbacks. The method requires that each sample has equal input amounts of the nucleic acid and that the amplification efficiency between samples is identical until the time of analysis.
10 Furthermore, it is difficult using the conventional methods of PCR quantitation such as gel electrophoresis or plate capture hybridization to determine that all samples are in fact analyzed during the log phase of the reaction as required by the method.

Another method called quantitative competitive (QC)-PCR, as the name implies, relies on the inclusion of an internal control competitor in each reaction [Piatak, (1993), BioTechniques]. The
15 efficiency of each reaction is normalized to the internal competitor. A known amount of internal competitor is typically added to each sample. The unknown target PCR product is compared with the known competitor PCR product to obtain relative quantitation. A difficulty with this general approach lies in developing an internal control that amplifies with the same efficiency than the target molecule.

20 *5' Fluorogenic Nuclease Assays*

Fluorogenic nuclease assays are a real time quantitation method that uses a probe to monitor formation of amplification product. The basis for this method of monitoring the formation of amplification product is to measure continuously PCR product accumulation using a dual-labelled fluorogenic oligonucleotide probe, an approach frequently referred to in the literature simply as the
25 "TaqMan method" [Piatak,(1993), Science; Heid, (1996); Gibson, (1996); Holland. (1991)].

The probe used in such assays is typically a short (about 20-25 bases) oligonucleotide that is labeled with two different fluorescent dyes. The 5' terminus of the probe is attached to a reporter dye and the 3' terminus is attached to a quenching dye, although the dyes could be attached at other locations on the probe as well. The probe is designed to have at least substantial sequence
30 complementarity with the probe binding site. Upstream and downstream PCR primers which bind to flanking regions of the locus are added to the reaction mixture. When the probe is intact, energy transfer between the two fluorophors occurs and the quencher quenches emission from the reporter. During the extension phase of PCR, the probe is cleaved by the 5' nuclease activity of a

nucleic acid polymerase such as Taq polymerase, thereby releasing the reporter from the oligonucleotide-quencher and resulting in an increase of reporter emission intensity which can be measured by an appropriate detector.

- One detector which is specifically adapted for measuring fluorescence emissions such as those created during a fluorogenic assay is the ABI 7700 or 4700 HT manufactured by Applied Biosystems, Inc. in Foster City, Calif. The ABI 7700 uses fiber optics connected with each well in a 96-or 384 well PCR tube arrangement. The instrument includes a laser for exciting the labels and is capable of measuring the fluorescence spectra intensity from each tube with continuous monitoring during PCR amplification. Each tube is re-examined every 8.5 seconds.
- Computer software provided with the instrument is capable of recording the fluorescence intensity of reporter and quencher over the course of the amplification. The recorded values will then be used to calculate the increase in normalized reporter emission intensity on a continuous basis. The increase in emission intensity is plotted versus time, i.e., the number of amplification cycles, to produce a continuous measure of amplification. To quantify the locus in each amplification reaction, the amplification plot is examined at a point during the log phase of product accumulation. This is accomplished by assigning a fluorescence threshold intensity above background and determining the point at which each amplification plot crosses the threshold (defined as the threshold cycle number or Ct). Differences in threshold cycle number are used to quantify the relative amount of PCR target contained within each tube. Assuming that each reaction functions at 100% PCR efficiency, a difference of one Ct represents a two-fold difference in the amount of starting template. The fluorescence value can be used in conjunction with a standard curve to determine the amount of amplification product present.

Non-Probe-Based Detection Methods

- A variety of options are available for measuring the amplification products as they are formed. One method utilizes labels, such as dyes, which only bind to double stranded DNA. In this type of approach, amplification product (which is double stranded) binds dye molecules in solution to form a complex. With the appropriate dyes, it is possible to distinguish between dye molecules free in solution and dye molecules bound to amplification product. For example, certain dyes fluoresce only when bound to amplification product. Examples of dyes which can be used in methods of this general type include, but are not limited to, Syber Green.TM. and Pico Green from Molecular Probes, Inc. of Eugene, Oreg., ethidium bromide, propidium iodide, chromomycin, acridine orange, Hoechst 33258, Toto-1, Yoyo-1, DAPI (4',6-diamidino-2-phenylindole hydrochloride).

Another real time detection technique measures alteration in energy fluorescence energy transfer between fluorophors conjugated with PCR primers [Livak, (1995)].

Probe-Based Detection Methods

5 These detection methods involve some alteration to the structure or conformation of a probe hybridized to the locus between the amplification primer pair. In some instances, the alteration is caused by the template-dependent extension catalyzed by a nucleic acid polymerase during the amplification process. The alteration generates a detectable signal which is an indirect measure of the amount of amplification product formed.

10 For example, some methods involve the degradation or digestion of the probe during the extension reaction. These methods are a consequence of the 5'-3' nuclease activity associated with some nucleic acid polymerases. Polymerases having this activity cleave mononucleotides or small oligonucleotides from an oligonucleotide probe annealed to its complementary sequence located within the locus.

15 The 3' end of the upstream primer provides the initial binding site for the nucleic acid polymerase. As the polymerase catalyzes extension of the upstream primer and encounters the bound probe, the nucleic acid polymerase displaces a portion of the 5' end of the probe and through its nuclease activity cleaves mononucleotides or oligonucleotides from the probe.

20 The upstream primer and the probe can be designed such that they anneal to the complementary strand in close proximity to one another. In fact, the 3' end of the upstream primer and the 5' end of the probe may abut one another. In this situation, extension of the upstream primer is not necessary in order for the nucleic acid polymerase to begin cleaving the probe. In the case in which intervening nucleotides separate the upstream primer and the probe, extension of the primer is necessary before the nucleic acid polymerase encounters the 5' end of the probe. Once contact occurs and polymerization continues, the 5'-3' exonuclease activity of the nucleic acid polymerase
25 begins cleaving mononucleotides or oligonucleotides from the 5' end of the probe. Digestion of the probe continues until the remaining portion of the probe dissociates from the complementary strand.

30 In solution, the two end sections can hybridize with each other to form a hairpin loop. In this conformation, the reporter and quencher dye are in sufficiently close proximity that fluorescence from the reporter dye is effectively quenched by the quencher dye. Hybridized probe, in contrast, results in a linearized conformation in which the extent of quenching is decreased. Thus, by moni-

toring emission changes for the two dyes, it is possible to indirectly monitor the formation of amplification product.

Probes

5 The labeled probe is selected so that its sequence is substantially complementary to a segment of the test locus or a reference locus. As indicated above, the nucleic acid site to which the probe binds should be located between the primer binding sites for the upstream and downstream amplification primers.

Primers

10 The primers used in the amplification are selected so as to be capable of hybridizing to sequences at flanking regions of the locus being amplified. The primers are chosen to have at least substantial complementarity with the different strands of the nucleic acid being amplified. When a probe is utilized to detect the formation of amplification products, the primers are selected in such that they flank the probe, i.e. are located upstream and downstream of the probe.

15 The primer must have sufficient length so that it is capable of priming the synthesis of extension products in the presence of an agent for polymerization. The length and composition of the primer depends on many parameters, including, for example, the temperature at which the annealing reaction is conducted, proximity of the probe binding site to that of the primer, relative concentrations of the primer and probe and the particular nucleic acid composition of the probe. Typically the primer includes 15-30 nucleotides. However, the length of the primer may be more
20 or less depending on the complexity of the primer binding site and the factors listed above.

Labels for Probes and Primers

The labels used for labeling the probes or primers of the current invention and which can provide the signal corresponding to the quantity of amplification product can take a variety of forms. As indicated above with regard to the 5' fluorogenic nuclease method, a fluorescent signal is one
25 signal which can be measured. However, measurements may also be made, for example, by monitoring radioactivity, colorimetry, absorption, magnetic parameters, or enzymatic activity. Thus, labels which can be employed include, but are not limited to, fluorophors, chromophores, radioactive isotopes, electron dense reagents, enzymes, and ligands having specific binding partners (e.g., biotin-avidin).

30 Monitoring changes in fluorescence is a particularly useful way to monitor the accumulation of amplification products. A number of labels useful for attachment to probes or primers are

commercially available including fluorescein and various fluorescein derivatives such as FAM, HEX, TET and JOE (all which are available from Applied Biosystems, Foster City, Calif.); lucifer yellow, and coumarin derivatives.

5 Labels may be attached to the probe or primer using a variety of techniques and can be attached at the 5' end, and/or the 3' end and/or at an internal nucleotide. The label can also be attached to spacer arms of various sizes which are attached to the probe or primer. These spacer arms are useful for obtaining a desired distance between multiple labels attached to the probe or primer.

10 In some instances, a single label may be utilized; whereas, in other instances, such as with the 5' fluorogenic nuclease assays for example; two or more labels are attached to the probe. In cases wherein the probe includes multiple labels, it is generally advisable to maintain spacing between the labels which is sufficient to permit separation of the labels during digestion of the probe through the 5'-3' nuclease activity of the nucleic acid polymerase.

Patients Exhibiting Symptoms of Disease

15 A number of diseases are associated with changes in the copy number of a certain gene. For patients having symptoms of a disease, the real-time PCR method can be used to determine if the patient has copy number alterations which are known to be linked with diseases that are associated with the symptoms the patient has.

GSK3B expression

GSK3B fusion proteins

20 Fusion proteins are useful for generating antibodies against GSK3B polypeptides and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of GSK3B polypeptides. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

25 A GSK3B fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment can comprise at least 54, 75, 100, 125, 139, 150, 175, 200, 225, 250, 275, 300, 325 or 350 contiguous amino acids of SEQ ID NO: 2 or of a biologically active variant, such as those described above. The first polypeptide segment also can comprise full-length GSK3B.

30 The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include, but are not limited to β galactosidase, β -

glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. A fusion protein also can be engineered to contain a cleavage site located adjacent to the GSK3B.

Preparation of Polynucleotides

10 A naturally occurring GSK3B polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique
15 for obtaining a polynucleotide can be used to obtain isolated GSK3B polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprise GSK3B nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

GSK3B cDNA molecules can be made with standard molecular biology techniques, using GSK3B
20 mRNA as a template. GSK3B cDNA molecules can thereafter be replicated using molecular biology techniques known in the art. An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

Alternatively, synthetic chemistry techniques can be used to synthesize GSK3B polynucleotides.
25 The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode GSK3B having, for example, an amino acid sequence shown in SEQ ID NO: 2 or a biologically active variant thereof.

Extending Polynucleotides

Various PCR-based methods can be used to extend nucleic acid sequences encoding human
30 GSK3B, for example to detect upstream sequences of GSK3B gene such as promoters and regulatory elements. For example, restriction-site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus. Genomic DNA is first amplified in the presence of a primer

to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

- 5 Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region. Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable
10 fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

- Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. In this method, multiple restriction enzyme digestions and ligations also can be used to place an
15 engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

- When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be
20 especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

- Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary
25 sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate equipment and software (*e.g.*, GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic
30 data display can be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

Obtaining Polypeptides

GSK3B can be obtained, for example, by purification from human cells, by expression of GSK3B polynucleotides, or by direct chemical synthesis.

Protein Purification

- 5 GSK3B can be purified from any human cell which expresses the enzyme, including those which have been transfected with expression constructs which express GSK3B. A purified GSK3B is separated from other compounds which normally associate with GSK3B in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange
10 chromatography, affinity chromatography, and preparative gel electrophoresis.

Expression of GSK3B Polynucleotides

- To express GSK3B, GSK3B polynucleotides can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct
15 expression vectors containing sequences encoding GSK3B and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination.

- A variety of expression vector/host systems can be utilized to contain and express sequences encoding GSK3B. These include, but are not limited to, microorganisms, such as bacteria trans-
20 formed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (*e.g.*, baculovirus), plant cell systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids), or animal cell systems.

- 25 The control elements or regulatory sequences are those non-translated regions of the vector - enhancers, promoters, 5' and 3' untranslated regions -- which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example,
30 when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells.

Promoters or enhancers derived from the genomes of plant cells (*e.g.*, heat shock, RUBISCO, and storage protein genes) or from plant viruses (*e.g.*, viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple
5 copies of a nucleotide sequence encoding GSK3B, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

Bacterial and Yeast Expression Systems

In bacterial systems, a number of expression vectors can be selected. For example, when a large quantity of GSK3B is needed for the induction of antibodies, vectors which direct high level
10 expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding GSK3B can be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced. pIN vectors or pGEX vectors (Promega,
15 Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be
20 released from the GST moiety at will.

Plant and Insect Expression Systems

If plant expression vectors are used, the expression of sequences encoding GSK3B can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV.
25 Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used. These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection.

An insect system also can be used to express GSK3B. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign
30 genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding GSK3B can be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of GSK3B will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can

then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which GSK3B can be expressed.

Mammalian Expression Systems

5 A number of viral-based expression systems can be used to express GSK3B in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding GSK3B can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing GSK3B in infected host cells [Engelhard, 1994]. If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (*e.g.*, liposomes, polycationic amino polymers, or vesicles). Specific initiation signals also can be used to achieve more efficient translation of sequences encoding GSK3B. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding GSK3B, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic.

Host Cells

25 A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed GSK3B in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipi-
dation, and acylation. Post-translational processing which cleaves a "prepro" form of the poly-
peptide also can be used to facilitate correct insertion, folding and/or function. Different host cells
which have specific cellular machinery and characteristic mechanisms for post-translational
30 activities (*e.g.*, CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express GSK3B can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the
5 vector, cells can be allowed to grow for 1-2 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced GSK3B sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. Any number of selection systems can be used to
10 recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase [Logan, (1984)] and adenine phosphoribosyltransferase [Wigler, (1977)] genes which can be employed in *tk⁻* or *aprt⁻* cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate [Lowy, (1980)], *npt* confers resistance to the aminoglycosides, neomycin and G-418
15 [Wigler, (1980)], and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively [Colbere-Garapin, 1981]. Additional selectable genes have been described. For example, *trpB* allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine. Visible markers such as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to
20 identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system

Detecting Polypeptide Expression

Although the presence of marker gene expression suggests that a GSK3B polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence
25 encoding GSK3B is inserted within a marker gene sequence, transformed cells containing sequences which encode GSK3B can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding GSK3B under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of GSK3B polynucleotide.

30 Alternatively, host cells which contain a GSK3B polynucleotide and which express GSK3B can be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a poly-

nucleotide sequence encoding GSK3B can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding GSK3B. Nucleic acid amplification-based assays involve the use of oligonucleotides selected from sequences encoding GSK3B to detect transformants which contain a GSK3B polynucleotide.

- 5 A variety of protocols for detecting and measuring the expression of GSK3B, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on GSK3B can be used, or a competitive
10 binding assay can be employed.

- A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding GSK3B include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide.
15 Alternatively, sequences encoding GSK3B can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes *in vitro* by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter
20 molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Expression and Purification of Polypeptides

- Host cells transformed with GSK3B polynucleotides can be cultured under conditions suitable for
25 the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing GSK3B polynucleotides can be designed to contain signal sequences which direct secretion of soluble GSK3B through a prokaryotic or eukaryotic cell membrane or which direct the membrane
30 insertion of membrane-bound GSK3B.

As discussed above, other constructions can be used to join a sequence encoding GSK3B to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating

peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and GSK3B also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing GSK3B and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography) Maddox, (1983)], while the enterokinase cleavage site provides a means for purifying GSK3B from the fusion protein [Porath, (1992)].

Chemical Synthesis

Sequences encoding GSK3B can be synthesized, in whole or in part, using chemical methods well known in the art. Alternatively, GSK3B itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques. Protein synthesis can either be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of GSK3B can be separately synthesized and combined using chemical methods to produce a full-length molecule.

The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography. The composition of a synthetic GSK3B can be confirmed by amino acid analysis or sequencing. Additionally, any portion of the amino acid sequence of GSK3B can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

Production of Altered Polypeptides

As will be understood by those of skill in the art, it may be advantageous to produce GSK3B polynucleotides possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

The nucleotide sequences referred to herein can be engineered using methods generally known in the art to alter GSK3B polynucleotides for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA

product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

5 *GSK3B* Analogs

One general class of GSK3B analogs are variants having an amino acid sequence that is a mutation of the amino acid sequence disclosed herein. Another general class of GSK3B analogs is provided by anti-idiotypic antibodies, and fragments thereof, as described below. Moreover, recombinant antibodies comprising anti-idiotypic variable domains can be used as analogs (see, for example, 10 [Monfardini et al., (1996)]). Since the variable domains of anti-idiotypic GSK3B antibodies mimic GSK3B, these domains can provide GSK3B enzymatic activity. Methods of producing anti-idiotypic catalytic antibodies are known to those of skill in the art [Joron et al., (1992), Friboulet et al. (1994), Avelle et al., (1998)].

Another approach to identifying GSK3B analogs is provided by the use of combinatorial libraries. 15 Methods for constructing and screening phage display and other combinatorial libraries are provided, for example, by [Kay et al., *Phage Display of Peptides and Proteins* (Academic Press 1996), U.S. 5,783,384, U.S. 5,747,334, and U.S. 5,723,323.

Antibodies

Any type of antibody known in the art can be generated to bind specifically to an epitope of 20 GSK3B.

"Antibody" as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab')₂, and Fv, which are capable of binding an epitope of GSK3B. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more; e.g., at least 15, 25, or 50 amino acid. An antibody which specifically binds to an epitope of GSK3B can be used 25 therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radio-immunoassays, immunohistochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are 30 well known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the GSK3B immunogen.

Typically, an antibody which specifically binds to GSK3B provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to GSK3B do not detect other proteins in immunochemical assays and can immunoprecipitate GSK3B from solution.

- 5 GSK3B can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, GSK3B can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (*e.g.*, aluminum
10 hydroxide), and surface active substances (*e.g.*, lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (*bacilli Calmette-Guerin*) and *Corynebacterium parvum* are especially useful.

Monoclonal antibodies which specifically bind to GSK3B can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These
15 techniques include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique [Roberge, (1995)].

In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. Monoclonal and other antibodies also can be
20 "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual
25 residues or by grating of entire complementarity determining regions. Antibodies which specifically bind to GSK3B can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. 5,565,332.

Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to
30 GSK3B. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries. Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template. Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught. A

nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology.

Antibodies which specifically bind to GSK3B also can be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents. Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared.

Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which GSK3B is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

Antisense Oligonucleotides

Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of GSK3B gene products in the cell.

Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters.

Modifications of GSK3B gene expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the GSK3B gene. Oligonucleotides derived from the transcription initiation site, *e.g.*, between positions -10 and +10 from

the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature [Nicholls, (1993)].

- 5 An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a GSK3B polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a GSK3B polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent GSK3B nucleotides, can provide sufficient targeting specificity for GSK3B mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular GSK3B polynucleotide sequence. Antisense oligonucleotides can be modified without affecting their ability to hybridize to a GSK3B polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art.

Ribozymes

Ribozymes are RNA molecules with catalytic activity [Uhlmann, (1987)]. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences. The coding sequence of a GSK3B polynucleotide can be used to generate ribozymes which will specifically bind to mRNA transcribed from a GSK3B polynucleotide. Methods of designing and constructing ribozymes

which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art. For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target RNA.

Specific ribozyme cleavage sites within a GSK3B RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate GSK3B RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. The nucleotide sequences shown in SEQ ID NO: 1 and its complement provide sources of suitable hybridization region sequences. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease GSK3B expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells (U.S. 5,641,673). Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

Screening / Screening Assays

30 *Regulators*

Regulators as used herein, refer to compounds that affect the activity of GSK3B in vivo and/or in vitro. Regulators can be agonists and antagonists of GSK3B polypeptide and can be compounds that exert their effect on the GSK3B activity via the enzymatic activity, expression, post-trans-

lational modifications or by other means. Agonists of GSK3B are molecules which, when bound to GSK3B, increase or prolong the activity of GSK3B. Agonists of GSK3B include proteins, nucleic acids, carbohydrates, small molecules, or any other molecule which activate GSK3B. Antagonists of GSK3B are molecules which, when bound to GSK3B, decrease the amount or the
5 duration of the activity of GSK3B. Antagonists include proteins, nucleic acids, carbohydrates, antibodies, small molecules, or any other molecule which decrease the activity of GSK3B.

The term "modulate", as it appears herein, refers to a change in the activity of GSK3B polypeptide. For example, modulation may cause an increase or a decrease in enzymatic activity, binding characteristics, or any other biological, functional, or immunological properties of
10 GSK3B.

As used herein, the terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein recognized by the binding molecule (i.e., the antigenic determinant or epitope). For example, if an antibody is specific for
15 epitope "A" the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The invention provides methods (also referred to herein as "screening assays") for identifying compounds which can be used for the treatment of diseases related to GSK3B. The methods entail
20 the identification of candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other molecules) which bind to GSK3B and/or have a stimulatory or inhibitory effect on the biological activity of GSK3B or its expression and then determining which of these compounds have an effect on symptoms or diseases related to GSK3B in an *in vivo* assay.

Candidate or test compounds or agents which bind to GSK3B and/or have a stimulatory or
25 inhibitory effect on the activity or the expression of GSK3B are identified either in assays that employ cells which express GSK3B (cell-based assays) or in assays with isolated GSK3B (cell-free assays). The various assays can employ a variety of variants of GSK3B (e.g., full-length GSK3B, a biologically active fragment of GSK3B, or a fusion protein which includes all or a portion of GSK3B). Moreover, GSK3B can be derived from any suitable mammalian species (e.g.,
30 human GSK3B, rat GSK3B or murine GSK3B). The assay can be a binding assay entailing direct or indirect measurement of the binding of a test compound or a known GSK3B ligand to GSK3B. The assay can also be an activity assay entailing direct or indirect measurement of the activity of GSK3B. The assay can also be an expression assay entailing direct or indirect measurement of the expression of GSK3B mRNA or GSK3B protein. The various screening assays are combined with

an *in vivo* assay entailing measuring the effect of the test compound on the symptoms of diseases related to GSK3B.

5 The present invention includes biochemical, cell free assays that allow the identification of inhibitors and agonists of kinases suitable as lead structures for pharmacological drug development. Such assays involve contacting a form of GSK3B (e.g., full-length GSK3B, a biologically active fragment of GSK3B, or a fusion protein comprising all or a portion of GSK3B) with a test compound and determining the ability of the test compound to act as an antagonist (preferably) or an agonist of the enzymatic activity of GSK3B.

10 The activity of GSK3B molecules of the present invention can be measured using a variety of assays that measure GSK3B activity. For example, GSK3B enzyme activity can be assessed by a standard *in vitro* kinase assay.

GSK3B can be used in substantial and specific assays related to kinases. Such assays involve any of the known kinase functions or activities or properties useful for diagnosis and treatment of kinase-related conditions that are specific for the subfamily of kinases that the one of the present
15 invention belongs to, particularly in cells and tissues that express the kinase.

GSK3B is also useful in drug screening assays, in cell-based or cell-free systems.

GSK3B can be used to identify compounds that modulate kinase activity of the protein in its natural state or an altered form that causes a specific disease or pathology associated with the kinase.

20 The present invention also relates to a method of screening compounds having inhibitory activity of kinase activity of the proteins of the present invention. This screening method consists of two steps. First, GSK3B is caused to contact a substrate to be phosphorylated by this protein in the presence of a test compound to detect the kinase activity of the protein of the present invention. Second, the kinase activity detected in step (a) is compared with that detected in the absence of the
25 test compound, and a compound that lowers the kinase activity of the protein of the present invention is selected.

The kinase activity of GSK3B can be detected, for example, by adding ATP having radioactively labeled phosphate to the reaction system containing the protein of the present invention and the substrate and measuring the radioactivity of the phosphate attached to the substrate.

30 Both, GSK3B and appropriate variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to modulate the kinase activity. These compounds can

be further screened against a functional kinase to determine the effect of the compound on the kinase activity. Further, these compounds can be tested in animal or invertebrate systems to determine activity/effectiveness. Compounds can be identified that activate (agonist) or inactivate (antagonist) the kinase to a desired degree.

- 5 Further, GSK3B can be used to screen a compound for the ability to stimulate or inhibit interaction between the kinase protein and a molecule that normally interacts with the kinase protein, e.g. a substrate or a component of the signal pathway that the kinase protein normally interacts (for example, another kinase). Such assays typically include the steps of combining the kinase protein with a candidate compound under conditions that allow the kinase protein, or fragment, to interact
10 with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the kinase protein and the target, such as any of the associated effects of signal transduction such as protein phosphorylation, cAMP turnover, and adenylate cyclase activation, etc.

- The invention further includes other end point assays to identify compounds that modulate
15 (stimulate or inhibit) GSK3B kinase activity. The assays typically involve an assay of events in the signal transduction pathway that indicate kinase activity. Thus, the phosphorylation of a substrate, activation of a protein, a change in the expression of genes that are up- or down-regulated in response to the kinase protein dependent signal cascade can be assayed.

- Any of the biological or biochemical functions mediated by the kinase can be used as an endpoint
20 assay. These include all of the biochemical or biochemical/biological events described herein, in the references cited herein, incorporated by reference for these endpoint assay targets, and other functions known to those of ordinary skill in the art.

- Binding and/or activating compounds can also be screened by using chimeric kinase proteins in which the amino terminal extracellular domain, or parts thereof, the entire transmembrane domain
25 or subregions, such as any of the seven transmembrane segments or any of the intracellular or extracellular loops and the carboxy terminal intracellular domain, or parts thereof, can be replaced by heterologous domains or subregions. For example, a substrate-binding region can be used that interacts with a different substrate than that which is recognized by the native kinase. Accordingly, a different set of signal transduction components is available as an end-point assay for activation.
30 This allows for assays to be performed in other than the specific host cell from which the kinase is derived.

GSK3B is also useful in competition binding assays in methods designed to discover compounds that interact with the kinase (e.g. binding partners and/or ligands). Thus, a compound is exposed to

a kinase polypeptide under conditions that allow the compound to bind or to otherwise interact with the polypeptide. Soluble kinase polypeptide is also added to the mixture. If the test compound interacts with the soluble kinase polypeptide, it decreases the amount of complex formed or activity from the kinase target. This type of assay is particularly useful in cases in which
5 compounds are sought that interact with specific regions of the kinase.

Agents that modulate GSK3B can be identified using one or more of the above assays, alone or in combination. It is generally preferable to use a cell-based or cell free system first and then confirm activity in an animal or other model system. Such model systems are well known in the art and can readily be employed in this context.

10 Test compounds used for this screening methods are not particularly limited and are generally low-molecular-weight compounds, proteins (including antibodies), peptides, etc. Test compounds are either artificially synthesized or natural.

As used herein, an antibody is defined in terms consistent with that recognized within the art: they are multi-subunit proteins produced by a mammalian organism in response to an antigen challenge.
15 The antibodies of the present invention include polyclonal antibodies and monoclonal antibodies, as well as fragments of such antibodies, including, but not limited to, Fab or F(ab').sub.2, and Fv fragments.

Many methods are known for generating and/or identifying antibodies to a given target peptide. Several such methods are described by [Harlow, (1989)].

20 In general, to generate antibodies, an isolated peptide is used as an immunogen and is administered to a mammalian organism, such as a rat, rabbit or mouse. The full-length protein, an antigenic peptide fragment or a fusion protein can be used. Particularly important fragments are those covering functional domains, such as the domains identified in FIG. 2, and domain of sequence homology or divergence amongst the family, such as those that can readily be identified using
25 protein alignment methods and as presented in the Figures.

Antibodies are preferably prepared from regions or discrete fragments of the kinase proteins. Antibodies can be prepared from any region of the peptide as described herein. However, preferred regions will include those involved in function/activity and/or kinase/binding partner interaction.

Modulators of kinase protein activity identified according to these drug screening assays can be
30 used to treat a subject with a disorder mediated by the kinase pathway, by treating cells or tissues that express the kinase.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a kinase-modulating agent, an antisense kinase nucleic acid molecule, a kinase-specific antibody, or a kinase-binding partner) can be used in an animal or other model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal or other model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

- 10 GSK3B proteins are also useful to provide a target for diagnosing a disease or predisposition to disease mediated by the peptide. Accordingly, the invention provides methods for detecting the presence, or levels of, the protein (or encoding mRNA) in a cell, tissue, or organism.

One agent for detecting a protein in a sample is an antibody capable of selectively binding to protein. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

- 20 The peptides of the present invention also provide targets for diagnosing active protein activity, disease, or predisposition to disease, in a patient having a variant peptide, particularly activities and conditions that are known for other members of the family of proteins to which the present one belongs. Thus, the peptide can be isolated from a biological sample and assayed for the presence of a genetic mutation that results in aberrant peptide. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered kinase activity in cell-based or cell-free assay, alteration in substrate or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein.

- 30 The diseases for which detection of kinase genes in a sample could be diagnostic include diseases in which kinase nucleic acid (DNA and/or RNA) is amplified in comparison to normal cells. By "amplification" is meant increased numbers of kinase DNA or RNA in a cell compared with normal cells. In normal cells, kinases are typically found as single copy genes. In selected diseases, the chromosomal location of the kinase genes may be amplified, resulting in multiple copies of the gene, or amplification. Gene amplification can lead to amplification of kinase RNA, or kinase RNA can be amplified in the absence of kinase DNA amplification.

In vitro techniques for detection of peptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence using a detection reagent, such as an antibody or protein binding agent. Alternatively, the peptide can be detected in vivo in a subject by introducing into the subject a labeled anti-peptide antibody or other types of
5 detection agent. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods that detect the allelic variant of a peptide expressed in a subject and methods which detect fragments of a peptide in a sample.

Solution in vitro assays can be used to identify a GSK3B substrate or inhibitor. Solid phase
10 systems can also be used to identify a substrate or inhibitor of a GSK3B polypeptide. For example, a GSK3B polypeptide or GSK3B fusion protein can be immobilized onto the surface of a receptor chip of a commercially available biosensor instrument (BIAcore, Biacore AB; Uppsala, Sweden). The use of this instrument is disclosed, for example, by [Karlsson, (1991), and Cunningham and Wells, (1993)].

15 In brief, a GSK3B polypeptide or fusion protein is covalently attached, using amine or sulfhydryl chemistry, to dextran fibers that are attached to gold film within a flow cell. A test sample is then passed through the cell. If a GSK3B substrate or inhibitor is present in the sample, it will bind to the immobilized polypeptide or fusion protein, causing a change in the refractive index of the medium, which is detected as a change in surface plasmon resonance of the gold film. This system
20 allows the determination on- and off-rates, from which binding affinity can be calculated, and assessment of the stoichiometry of binding, as well as the kinetic effects of GSK3B mutation. This system can also be used to examine antibody-antigen interactions, and the interactions of other complement/anti-complement pairs.

In one embodiment, the invention provides assays for screening candidate or test compounds
25 which bind to or modulate the activity of GSK3B. Such assays can employ full-length GSK3B, a biologically active fragment of GSK3B, or a fusion protein which includes all or a portion of GSK3B. As described in greater detail below, the test compound can be obtained by any suitable means, e.g., from conventional compound libraries.

Determining the ability of the test compound to modulate the activity of GSK3B can be
30 accomplished, for example, by determining the ability of GSK3B to bind to or interact with a target molecule. The target molecule can be a molecule with which GSK3B binds or interacts with in nature. The target molecule can be a component of a signal transduction pathway which facilitates transduction of an extracellular signal. The target GSK3B molecule can be, for example,

a second intracellular protein which has catalytic activity or a protein which facilitates the association of downstream signaling molecules with GSK3B.

Determining the ability of GSK3B to bind to or interact with a target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of a polypeptide of the invention to bind to or interact with a target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (e.g., intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (e.g., a regulatory element that is responsive to a polypeptide of the invention operably linked to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a cellular response.

In various embodiments of the above assay methods of the present invention, it may be desirable to immobilize GSK3B (or a GSK3B target molecule) to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to GSK3B, or interaction of GSK3B with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and microcentrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase (GST) fusion proteins or glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical; St. Louis, Mo.) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or GSK3B, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components and complex formation is measured either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of binding or activity of GSK3B can be determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either GSK3B or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated polypeptide of the invention or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, Ill.), and immobilized in the wells of

streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies reactive with GSK3B or target molecules but which do not interfere with binding of the polypeptide of the invention to its target molecule can be derivatized to the wells of the plate, and unbound target or polypeptide of the invention trapped in the wells by antibody conjugation. Methods for detecting such
5 complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with GSK3B or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with GSK3B or target molecule.

Another technique for drug screening which may be used provides for high throughput screening
10 of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with GSK3B, or fragments thereof, and washed. Bound GSK3B is then detected by methods well known in the art. Purified GSK3B can also be coated directly onto
15 plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding GSK3B specifically compete with a test compound for binding GSK3B. In this manner, antibodies can be used to detect the presence of any peptide which shares
20 one or more antigenic determinants with GSK3B.

The screening assay can also involve monitoring the expression of GSK3B. For example, regulators of expression of GSK3B can be identified in a method in which a cell is contacted with a candidate compound and the expression of GSK3B protein or mRNA in the cell is determined. The level of expression of GSK3B protein or mRNA the presence of the candidate compound is
25 compared to the level of expression of GSK3B protein or mRNA in the absence of the candidate compound. The candidate compound can then be identified as a regulator of expression of GSK3B based on this comparison. For example, when expression of GSK3B protein or mRNA protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of GSK3B protein or mRNA
30 expression. Alternatively, when expression of GSK3B protein or mRNA is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of GSK3B protein or mRNA expression. The level of GSK3B protein or mRNA expression in the cells can be determined by methods described below.

Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses GSK3B polynucleotide can be used in a cell-based assay system. The GSK3B polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line can be used.

5 *Binding Assays*

For binding assays, the test compound is preferably a small molecule which binds to and occupies the active site of GSK3B polypeptide, thereby making the ligand binding site inaccessible to substrate such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules. Potential ligands which
10 bind to a polypeptide of the invention include, but are not limited to, the natural ligands of known GSK3B kinases and analogues or derivatives thereof.

In binding assays, either the test compound or the GSK3B polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is
15 bound to GSK3B polypeptide can then be accomplished, for example, by direct counting of radioemmission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product. Alternatively, binding of a test compound to a GSK3B polypeptide can be determined without labeling either of the interactants. For example, a microphysio-
20 meter can be used to detect binding of a test compound with a GSK3B polypeptide. A microphysiometer (*e.g.*, Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and GSK3B [Haseloff, (1988)].

Determining the ability of a test compound to bind to GSK3B also can be accomplished using a
25 technology such as real-time Bimolecular Interaction Analysis (BIA) [McConnell, (1992); Sjolander, (1991)]. BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (*e.g.*, BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

30 In yet another aspect of the invention, a GSK3B-like polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay [Szabo, (1995); U.S. 5,283,317], to identify other proteins which bind to or interact with GSK3B and modulate its activity.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding GSK3B can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4).

- 5 In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact *in vivo* to form a protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ),
10 which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with GSK3B.

- It may be desirable to immobilize either the GSK3B (or polynucleotide) or the test compound to
15 facilitate separation of the bound form from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the GSK3B-like polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass
20 beads). Any method known in the art can be used to attach GSK3B-like polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked.
25 Binding of a test compound to GSK3B (or a polynucleotide encoding for GSK3B) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

- In one embodiment, GSK3B is a fusion protein comprising a domain that allows binding of GSK3B to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed
30 onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the non-adsorbed GSK3B; the mixture is then incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants

can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either GSK3B (or a polynucleotide encoding GSK3B) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated GSK3B (or a polynucleotide encoding biotinylated GSK3B) or test compounds can be prepared from biotin-NHS (N-hydroxysuccinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies which specifically bind to GSK3B, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the active site of GSK3B, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to GSK3B polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of GSK3B polypeptide, and SDS gel electrophoresis under non-reducing conditions.

Screening for test compounds which bind to a GSK3B polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a GSK3B polypeptide or polynucleotide can be used in a cell-based assay system. A GSK3B polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to GSK3B or a polynucleotide encoding GSK3B is determined as described above.

Functional Assays

Test compounds can be tested for the ability to increase or decrease GSK3B activity of a GSK3B polypeptide. The GSK3B activity can be measured, for example, using methods described in the specific examples, below. GSK3B activity can be measured after contacting either a purified GSK3B or an intact cell with a test compound. A test compound which decreases GSK3B activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for decreasing GSK3B activity. A test compound which increases GSK3B activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for increasing GSK3B activity.

Gene Expression

In another embodiment, test compounds which increase or decrease GSK3B gene expression are identified. As used herein, the term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding GSK3B, by northern analysis or realtime PCR is indicative of the presence of nucleic acids encoding GSK3B in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding GSK3B. The term "microarray", as used herein, refers to an array of distinct polynucleotides or oligonucleotides arrayed on a substrate, such as paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support. A GSK3B polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of GSK3B polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a regulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

The level of GSK3B mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of GSK3B polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined *in vivo*, in a cell culture, or in an *in vitro* translation system by detecting incorporation of labelled amino acids into GSK3B.

Test Compounds

Suitable test compounds for use in the screening assays of the invention can be obtained from any suitable source, e.g., conventional compound libraries. The test compounds can also be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds [Lam, (1997)]. Examples of

methods for the synthesis of molecular libraries can be found in the art. Libraries of compounds may be presented in solution or on beads, bacteria, spores, plasmids or phage.

Modeling of Regulators

Computer modeling and searching technologies permit identification of compounds, or the
5 improvement of already identified compounds, that can modulate GSK3B expression or activity. Having identified such a compound or composition, the active sites or regions are identified. Such sites might typically be the enzymatic active site, regulator binding sites, or ligand binding sites. The active site can be identified using methods known in the art including, for example, from the amino acid sequences of peptides, from the nucleotide sequences of nucleic acids, or from study of
10 complexes of the relevant compound or composition with its natural ligand. In the latter case, chemical or X-ray crystallographic methods can be used to find the active site by finding where on the factor the complexed ligand is found.

Next, the three dimensional geometric structure of the active site is determined. This can be done by known methods, including X-ray crystallography, which can determine a complete molecular
15 structure. On the other hand, solid or liquid phase NMR can be used to determine certain intramolecular distances. Any other experimental method of structure determination can be used to obtain partial or complete geometric structures. The geometric structures may be measured with a complexed ligand, natural or artificial, which may increase the accuracy of the active site structure determined.

20 If an incomplete or insufficiently accurate structure is determined, the methods of computer based numerical modeling can be used to complete the structure or improve its accuracy. Any recognized modeling method may be used, including parameterized models specific to particular biopolymers such as proteins or nucleic acids, molecular dynamics models based on computing molecular motions, statistical mechanics models based on thermal ensembles, or combined models.
25 For most types of models, standard molecular force fields, representing the forces between constituent atoms and groups, are necessary, and can be selected from force fields known in physical chemistry. The incomplete or less accurate experimental structures can serve as constraints on the complete and more accurate structures computed by these modeling methods.

Finally, having determined the structure of the active site, either experimentally, by modeling, or
30 by a combination, candidate modulating compounds can be identified by searching databases containing compounds along with information on their molecular structure. Such a search seeks compounds having structures that match the determined active site structure and that interact with

the groups defining the active site. Such a search can be manual, but is preferably computer assisted. These compounds found from this search are potential GSK3B modulating compounds.

Alternatively, these methods can be used to identify improved modulating compounds from an already known modulating compound or ligand. The composition of the known compound can be modified and the structural effects of modification can be determined using the experimental and computer modeling methods described above applied to the new composition. The altered structure is then compared to the active site structure of the compound to determine if an improved fit or interaction results. In this manner systematic variations in composition, such as by varying side groups, can be quickly evaluated to obtain modified modulating compounds or ligands of improved specificity or activity.

Therapeutic Indications and Methods

It was found by the present applicant that GSK3B is expressed in various human tissues.

Neurology

CNS disorders include disorders of the central nervous system as well as disorders of the peripheral nervous system.

CNS disorders include, but are not limited to brain injuries, cerebrovascular diseases and their consequences, Parkinson's disease, corticobasal degeneration, motor neuron disease, dementia, including ALS, multiple sclerosis, traumatic brain injury, stroke, post-stroke, post-traumatic brain injury, and small-vessel cerebrovascular disease. Dementias, such as Alzheimer's disease, vascular dementia, dementia with Lewy bodies, frontotemporal dementia and Parkinsonism linked to chromosome 17, frontotemporal dementias, including Pick's disease, progressive nuclear palsy, corticobasal degeneration, Huntington's disease, thalamic degeneration, Creutzfeld-Jakob dementia, HIV dementia, schizophrenia with dementia, and Korsakoff's psychosis, within the meaning of the definition are also considered to be CNS disorders.

Similarly, cognitive-related disorders, such as mild cognitive impairment, age-associated memory impairment, age-related cognitive decline, vascular cognitive impairment, attention deficit disorders, attention deficit hyperactivity disorders, and memory disturbances in children with learning disabilities are also considered to be CNS disorders.

Pain, within the meaning of this definition, is also considered to be a CNS disorder. Pain can be associated with CNS disorders, such as multiple sclerosis, spinal cord injury, sciatica, failed back surgery syndrome, traumatic brain injury, epilepsy, Parkinson's disease, post-stroke, and vascular

lesions in the brain and spinal cord (e.g., infarct, hemorrhage, vascular malformation). Non-central neuropathic pain includes that associated with post mastectomy pain, phantom feeling, reflex sympathetic dystrophy (RSD), trigeminal neuralgia, radioculopathy, post-surgical pain, HIV/AIDS related pain, cancer pain, metabolic neuropathies (e.g., diabetic neuropathy, vasculitic neuropathy secondary to connective tissue disease), paraneoplastic polyneuropathy associated, for example, with carcinoma of lung, or leukemia, or lymphoma, or carcinoma of prostate, colon or stomach, trigeminal neuralgia, cranial neuralgias, and post-herpetic neuralgia. Pain associated with peripheral nerve damage, central pain (i.e. due to cerebral ischemia) and various chronic pain i.e., lumbago, back pain (low back pain), inflammatory and/or rheumatic pain. Headache pain (for example, migraine with aura, migraine without aura, and other migraine disorders), episodic and chronic tension-type headache, tension-type like headache, cluster headache, and chronic paroxysmal hemicrania are also CNS disorders.

Visceral pain such as pancreatitis, intestinal cystitis, dysmenorrhea, irritable Bowel syndrome, Crohn's disease, biliary colic, ureteral colic, myocardial infarction and pain syndromes of the pelvic cavity, e.g., vulvodynia, orchialgia, urethral syndrome and protatodynia are also CNS disorders.

Also considered to be a disorder of the nervous system are acute pain, for example postoperative pain, and pain after trauma.

The human GSK3B is highly expressed in the following brain tissues: brain, cerebral cortex, frontal lobe, occipital lobe, parietal lobe, temporal lobe, caudatum, nucleus accumbens, putamen, posteroventral thalamus, dorsomedial thalamus, hypothalamus, spinal cord (ventral horn), spinal cord (dorsal horn), astrocytes. The expression in brain tissues demonstrates that the human GSK3B or mRNA can be utilized to diagnose nervous system diseases. Additionally the activity of the human GSK3B can be modulated to treat nervous system diseases.

25 *Cardiovascular Disorders*

Heart failure is defined as a pathophysiological state in which an abnormality of cardiac function is responsible for the failure of the heart to pump blood at a rate commensurate with the requirement of the metabolizing tissue. It includes all forms of pumping failures such as high-output and low-output, acute and chronic, right-sided or left-sided, systolic or diastolic, independent of the underlying cause.

Myocardial infarction (MI) is generally caused by an abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by arteriosclerosis. MI

prophylaxis (primary and secondary prevention) is included as well as the acute treatment of MI and the prevention of complications.

Ischemic diseases are conditions in which the coronary flow is restricted resulting in a perfusion which is inadequate to meet the myocardial requirement for oxygen. This group of diseases
5 includes stable angina, unstable angina and asymptomatic ischemia.

Arrhythmias include all forms of atrial and ventricular tachyarrhythmias, atrial tachycardia, atrial flutter, atrial fibrillation, atrio-ventricular reentrant tachycardia, preexcitation syndrome, ventricular tachycardia, ventricular flutter, ventricular fibrillation, as well as bradycardic forms of
10 arrhythmias.

Hypertensive vascular diseases include primary as well as all kinds of secondary arterial hypertension, renal, endocrine, neurogenic, others. The genes may be used as drug targets for the treatment of hypertension as well as for the prevention of all complications arising from cardiovascular diseases.

15 Peripheral vascular diseases are defined as vascular diseases in which arterial and/or venous flow is reduced resulting in an imbalance between blood supply and tissue oxygen demand. It includes chronic peripheral arterial occlusive disease (PAOD), acute arterial thrombosis and embolism, inflammatory vascular disorders, Raynaud's phenomenon and venous disorders.

Atherosclerosis is a cardiovascular disease in which the vessel wall is remodeled, compromising
20 the lumen of the vessel. The atherosclerotic remodeling process involves accumulation of cells, both smooth muscle cells and monocyte/macrophage inflammatory cells, in the intima of the vessel wall. These cells take up lipid, likely from the circulation, to form a mature atherosclerotic lesion. Although the formation of these lesions is a chronic process, occurring over decades of an adult human life, the majority of the morbidity associated with atherosclerosis occurs when a lesion
25 ruptures, releasing thrombogenic debris that rapidly occludes the artery. When such an acute event occurs in the coronary artery, myocardial infarction can ensue, and in the worst case, can result in death.

The formation of the atherosclerotic lesion can be considered to occur in five overlapping stages such as migration, lipid accumulation, recruitment of inflammatory cells, proliferation of vascular
30 smooth muscle cells, and extracellular matrix deposition. Each of these processes can be shown to occur in man and in animal models of atherosclerosis, but the relative contribution of each to the pathology and clinical significance of the lesion is unclear.

Thus, a need exists for therapeutic methods and agents to treat cardiovascular pathologies, such as atherosclerosis and other conditions related to coronary artery disease.

Cardiovascular diseases include but are not limited to disorders of the heart and the vascular system like congestive heart failure, myocardial infarction, ischemic diseases of the heart, all
5 kinds of atrial and ventricular arrhythmias, hypertensive vascular diseases, peripheral vascular diseases, and atherosclerosis.

Too high or too low levels of fats in the bloodstream, especially cholesterol, can cause long-term problems. The risk to develop atherosclerosis and coronary artery or carotid artery disease (and thus the risk of having a heart attack or stroke) increases with the total cholesterol level increasing.
10 Nevertheless, extremely low cholesterol levels may not be healthy. Examples of disorders of lipid metabolism are hyperlipidemia (abnormally high levels of fats (cholesterol, triglycerides, or both) in the blood, may be caused by family history of hyperlipidemia), obesity, a high-fat diet, lack of exercise, moderate to high alcohol consumption, cigarette smoking, poorly controlled diabetes, and an underactive thyroid gland), hereditary hyperlipidemias (type I hyperlipoproteinemia (familial
15 hyperchylomicronemia), type II hyperlipoproteinemia (familial hypercholesterolemia), type III hyperlipoproteinemia, type IV hyperlipoproteinemia, or type V hyperlipoproteinemia), hypolipoproteinemia, lipidoses (caused by abnormalities in the enzymes that metabolize fats), Gaucher's disease, Niemann-Pick disease, Fabry's disease, Wolman's disease, cerebrotendinous xanthomatosis, sitosterolemia, Refsum's disease, or Tay-Sachs disease.

20 Kidney disorders may lead to hypertension or hypotension. Examples for kidney problems possibly leading to hypertension are renal artery stenosis, pyelonephritis, glomerulonephritis, kidney tumors, polycystic kidney disease, injury to the kidney, or radiation therapy affecting the kidney. Excessive urination may lead to hypotension.

The human GSK3B is highly expressed in the following cardiovascular related tissues: heart
25 myocardial infarction, heart myocardial infarction, heart myocardial infarction, pericardium, heart atrium (right), heart atrium (left), heart ventricle (left), heart ventricle (right), Purkinje fibers, interventricular septum, aorta valve, coronary artery endothel cells, aortic smooth muscle cells, aortic endothel cells, liver tumor, adipose, fetal kidney, kidney, kidney, kidney tumor, renal epithelial cells, HEK 293 cells. Expression in the above mentioned tissues and in particular the
30 differential expression between diseased tissue heart myocardial infarction and healthy tissue demonstrates that the human GSK3B or mRNA can be utilized to diagnose of cardiovascular diseases. Additionally the activity of the human GSK3B can be modulated to treat cardiovascular diseases.

The human GSK3B is highly expressed in liver tissues: liver tumor. Expression in liver tissues demonstrates that the human GSK3B or mRNA can be utilized to diagnose of dyslipidemia disorders as an cardiovascular disorder. Additionally the activity of the human GSK3B can be modulated to treat - but not limited to - dyslipidemia disorders.

- 5 The human GSK3B is highly expressed in adipose tissues. Expression in adipose demonstrates that the human GSK3B or mRNA can be utilized to diagnose of dyslipidemia diseases as an cardiovascular disorder. Additionally the activity of the human GSK3B can be modulated to treat - but not limited to - dyslipidemia diseases.

- 10 The human GSK3B is highly expressed in kidney tissues : fetal kidney, kidney, kidney, kidney tumor, HEK 293 cells. Expression in kidney tissues demonstrates that the human GSK3B or mRNA can be utilized to diagnose of blood pressure disorders as an cardiovascular disorder. Additionally the activity of the human GSK3B can be modulated to treat - but not limited to - blood pressure disorders as hypertension or hypotension.

Hematological Disorders

- 15 Hematological disorders comprise diseases of the blood and all its constituents as well as diseases of organs and tissues involved in the generation or degradation of all the constituents of the blood. They include but are not limited to 1) Anemias, 2) Myeloproliferative Disorders, 3) Hemorrhagic Disorders, 4) Leukopenia, 5) Eosinophilic Disorders, 6) Leukemias, 7) Lymphomas, 8) Plasma Cell Dyscrasias, 9) Disorders of the Spleen in the course of hematological disorders. Disorders
20 according to 1) include, but are not limited to anemias due to defective or deficient hem synthesis, deficient erythropoiesis. Disorders according to 2) include, but are not limited to polycythemia vera, tumor-associated erythrocytosis, myelofibrosis, thrombocythemia. Disorders according to 3) include, but are not limited to vasculitis, thrombocytopenia, heparin-induced thrombocytopenia, thrombotic thrombocytopenic purpura, hemolytic-uremic syndrome, hereditary and acquired
25 disorders of platelet function, hereditary coagulation disorders. Disorders according to 4) include, but are not limited to neutropenia, lymphocytopenia. Disorders according to 5) include, but are not limited to hypereosinophilia, idiopathic hypereosinophilic syndrome. Disorders according to 6) include, but are not limited to acute myeloid leukemia, acute lymphoblastic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia, myelodysplastic syndrome. Disorders
30 according to 7) include, but are not limited to Hodgkin's disease, non-Hodgkin's lymphoma, Burkitt's lymphoma, mycosis fungoides cutaneous T-cell lymphoma. Disorders according to 8) include, but are not limited to multiple myeloma, macroglobulinemia, heavy chain diseases. In extension of the preceding idiopathic thrombocytopenic purpura, iron deficiency anemia, megaloblastic anemia (vitamin B12 deficiency), aplastic anemia, thalassemia, malignant

lymphoma bone marrow invasion, malignant lymphoma skin invasion, hemolytic uremic syndrome, giant platelet disease are considered to be hematological diseases too.

The human GSK3B is highly expressed in the following tissues of the hematological system: Jurkat (T-cells), Raji (B-cells), neutrophils cord blood, T-cells peripheral blood CD8+, monocytes
 5 peripheral blood CD14+, neutrophils peripheral blood, spleen. The expression in the above mentioned tissues demonstrates that the human GSK3B or mRNA can be utilized to diagnose of hematological diseases. Additionally the activity of the human GSK3B can be modulated to treat
 . hematological disorders.

Cancer Disorders

10 Cancer disorders within the scope of this definition comprise any disease of an organ or tissue in mammals characterized by poorly controlled or uncontrolled multiplication of normal or abnormal cells in that tissue and its effect on the body as a whole. Cancer diseases within the scope of the definition comprise benign neoplasms, dysplasias, hyperplasias as well as neoplasms showing
 15 metastatic growth or any other transformations like e.g. leukoplakias which often precede a breakout of cancer. Cells and tissues are cancerous when they grow more rapidly than normal cells, displacing or spreading into the surrounding healthy tissue or any other tissues of the body described as metastatic growth, assume abnormal shapes and sizes, show changes in their nucleocytoplasmatic ratio, nuclear polychromasia, and finally may cease. Cancerous cells and tissues may affect the body as a whole when causing paraneoplastic syndromes or if cancer occurs within
 20 a vital organ or tissue, normal function will be impaired or halted, with possible fatal results. The ultimate involvement of a vital organ by cancer, either primary or metastatic, may lead to the death of the mammal affected. Cancer tends to spread, and the extent of its spread is usually related to an individual's chances of surviving the disease. Cancers are generally said to be in one of three stages of growth: early, or localized, when a tumor is still confined to the tissue of origin, or
 25 primary site; direct extension, where cancer cells from the tumour have invaded adjacent tissue or have spread only to regional lymph nodes; or metastasis, in which cancer cells have migrated to distant parts of the body from the primary site, via the blood or lymph systems, and have established secondary sites of infection. Cancer is said to be malignant because of its tendency to cause death if not treated. Benign tumors usually do not cause death, although they may if they
 30 interfere with a normal body function by virtue of their location, size, or paraneoplastic side effects. Hence benign tumors fall under the definition of cancer within the scope of this definition as well. In general, cancer cells divide at a higher rate than do normal cells, but the distinction between the growth of cancerous and normal tissues is not so much the rapidity of cell division in the former as it is the partial or complete loss of growth restraint in cancer cells and their failure to

differentiate into a useful, limited tissue of the type that characterizes the functional equilibrium of growth of normal tissue. Cancer tissues may express certain molecular receptors and probably are influenced by the host's susceptibility and immunity and it is known that certain cancers of the breast and prostate, for example, are considered dependent on specific hormones for their existence. The term "cancer" under the scope of the definition is not limited to simple benign neoplasia but comprises any other benign and malign neoplasia like 1) Carcinoma, 2) Sarcoma, 3) Carcinosarcoma, 4) Cancers of the blood-forming tissues, 5) tumors of nerve tissues including the brain, 6) cancer of skin cells. Cancer according to 1) occurs in epithelial tissues, which cover the outer body (the skin) and line mucous membranes and the inner cavitory structures of organs e.g. such as the breast, lung, the respiratory and gastrointestinal tracts, the endocrine glands, and the genitourinary system. Ductal or glandular elements may persist in epithelial tumors, as in adenocarcinomas like e.g. thyroid adenocarcinoma, gastric adenocarcinoma, uterine adenocarcinoma. Cancers of the pavement-cell epithelium of the skin and of certain mucous membranes, such as e.g. cancers of the tongue, lip, larynx, urinary bladder, uterine cervix, or penis, may be termed epidermoid or squamous-cell carcinomas of the respective tissues and are in the scope of the definition of cancer as well. Cancer according to 2) develops in connective tissues, including fibrous tissues, adipose (fat) tissues, muscle, blood vessels, bone, and cartilage like e.g. osteogenic sarcoma; liposarcoma, fibrosarcoma, synovial sarcoma. Cancer according to 3) is cancer that develops in both epithelial and connective tissue. Cancer disease within the scope of this definition may be primary or secondary, whereby primary indicates that the cancer originated in the tissue where it is found rather than was established as a secondary site through metastasis from another lesion. Cancers and tumor diseases within the scope of this definition may be benign or malign and may affect all anatomical structures of the body of a mammal. By example but not limited to they comprise cancers and tumor diseases of I) the bone marrow and bone marrow derived cells (leukemias), II) the endocrine and exocrine glands like e.g. thyroid, parathyroid, pituitary, adrenal glands, salivary glands, pancreas III) the breast, like e.g. benign or malignant tumors in the mammary glands of either a male or a female, the mammary ducts, adenocarcinoma, medullary carcinoma, comedo carcinoma, Paget's disease of the nipple, inflammatory carcinoma of the young woman, IV) the lung, V) the stomach, VI) the liver and spleen, VII) the small intestine, VIII) the colon, IX) the bone and its supportive and connective tissues like malignant or benign bone tumour, e.g. malignant osteogenic sarcoma, benign osteoma, cartilage tumors; like malignant chondrosarcoma or benign chondroma; bone marrow tumors like malignant myeloma or benign eosinophilic granuloma, as well as metastatic tumors from bone tissues at other locations of the body; X) the mouth, throat, larynx, and the esophagus, XI) the urinary bladder and the internal and external organs and structures of the urogenital system of male and female like ovaries, uterus, cervix of the uterus, testes, and prostate gland, XII) the prostate, XIII) the pancreas, like ductal

carcinoma of the pancreas; XIV) the lymphatic tissue like lymphomas and other tumors of lymphoid origin, XV) the skin, XVI) cancers and tumor diseases of all anatomical structures belonging to the respiration and respiratory systems including thoracal muscles and linings, XVII) primary or secondary cancer of the lymph nodes XVIII) the tongue and of the bony structures of the hard palate or sinuses, XXIV) the mouth, cheeks, neck and salivary glands, XX) the blood vessels including the heart and their linings, XXI) the smooth or skeletal muscles and their ligaments and linings, XXII) the peripheral, the autonomous, the central nervous system including the cerebellum, XXIII) the adipose tissue.

The human GSK3B is highly expressed in the following cancer tissues: thyroid tumor, esophagus tumor, rectum tumor, liver tumor, Jurkat (T-cells), Raji (B-cells), lung tumor, ovary tumor, kidney tumor, HEK 293 cells. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue thyroid tumor and healthy tissue thyroid, between diseased tissue esophagus tumor and healthy tissue esophagus, between diseased tissue rectum tumor and healthy tissue rectum, between diseased tissue liver tumor and healthy tissue liver, between diseased tissue Jurkat (T-cells) and healthy tissue T-cells peripheral blood CD4+, between diseased tissue Raji (B-cells) and healthy tissue B-cells peripheral blood CD19+, between diseased tissue lung tumor and healthy tissue lung, between diseased tissue ovary tumor and healthy tissue ovary, between diseased tissue kidney tumor and healthy tissue kidney, between diseased tissue HEK 293 cells and healthy tissue kidney demonstrates that the human GSK3B or mRNA can be utilized to diagnose of cancer. Additionally the activity of the human GSK3B can be modulated to treat cancer.

Inflammatory Diseases

Inflammatory diseases comprise diseases triggered by cellular or non-cellular mediators of the immune system or tissues causing the inflammation of body tissues and subsequently producing an acute or chronic inflammatory condition. Examples for such inflammatory diseases are hypersensitivity reactions of type I – IV, for example but not limited to hypersensitivity diseases of the lung including asthma, atopic diseases, allergic rhinitis or conjunctivitis, angioedema of the lids, hereditary angioedema, antireceptor hypersensitivity reactions and autoimmune diseases, Hashimoto's thyroiditis, systemic lupus erythematosus, Goodpasture's syndrome, pemphigus, myasthenia gravis, Grave's and Raynaud's disease, type B insulin-resistant diabetes, rheumatoid arthritis, psoriasis, Crohn's disease, scleroderma, mixed connective tissue disease, polymyositis, sarcoidosis, glomerulonephritis, acute or chronic host versus graft reactions.

The human GSK3B is highly expressed in the following tissues of the immune system and tissues responsive to components of the immune system as well as in the following tissues responsive to

mediators of inflammation: neutrophils cord blood, neutrophils peripheral blood. The expression in the above mentioned tissues demonstrates that the human GSK3B or mRNA can be utilized to diagnose of inflammatory diseases. Additionally the activity of the human GSK3B can be modulated to treat inflammatory diseases.

5 *Disorders Related to Pulmology*

Asthma is thought to arise as a result of interactions between multiple genetic and environmental factors and is characterized by three major features: 1) intermittent and reversible airway obstruction caused by bronchoconstriction, increased mucus production, and thickening of the walls of the airways that leads to a narrowing of the airways, 2) airway hyperresponsiveness, and 10 3) airway inflammation. Certain cells are critical to the inflammatory reaction of asthma and they include T cells and antigen presenting cells, B cells that produce IgE, and mast cells, basophils, eosinophils, and other cells that bind IgE. These effector cells accumulate at the site of allergic reaction in the airways and release toxic products that contribute to the acute pathology and eventually to tissue destruction related to the disorder. Other resident cells, such as smooth muscle 15 cells, lung epithelial cells, mucus-producing cells, and nerve cells may also be abnormal in individuals with asthma and may contribute to its pathology. While the airway obstruction of asthma, presenting clinically as an intermittent wheeze and shortness of breath, is generally the most pressing symptom of the disease requiring immediate treatment, the inflammation and tissue destruction associated with the disease can lead to irreversible changes that eventually make 20 asthma a chronic and disabling disorder requiring long-term management.

Chronic obstructive pulmonary (or airways) disease (COPD) is a condition defined physiologically as airflow obstruction that generally results from a mixture of emphysema and peripheral airway obstruction due to chronic bronchitis [Botstein, 1980]. Emphysema is characterised by destruction of alveolar walls leading to abnormal enlargement of the air spaces of the lung. Chronic bronchitis 25 is defined clinically as the presence of chronic productive cough for three months in each of two successive years. In COPD, airflow obstruction is usually progressive and is only partially reversible. By far the most important risk factor for development of COPD is cigarette smoking, although the disease does also occur in non-smokers.

The human GSK3B is highly expressed in the following tissues of the respiratory system: 30 neutrophils cord blood, neutrophils peripheral blood, fetal lung, fetal lung fibroblast IMR-90 cells, fetal lung fibroblast MRC-5 cells, lung tumor, primary bronchia, bronchial epithelial cells, bronchial smooth muscle cells, small airway epithelial cells. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue fetal lung fibroblast IMR-90 cells and healthy tissue fetal lung, between diseased tissue fetal lung fibroblast

MRC-5 cells and healthy tissue fetal lung demonstrates that the human GSK3B or mRNA can be utilized to diagnose of respiratory diseases. Additionally the activity of the human GSK3B can be modulated to treat those diseases.

Disorders Related to Urology

- 5 Genitourinary disorders comprise benign and malign disorders of the organs constituting the genitourinary system of female and male, renal diseases like acute or chronic renal failure, immunologically mediated renal diseases like renal transplant rejection, lupus nephritis, immune complex renal diseases, glomerulopathies, nephritis, toxic nephropathy, obstructive uropathies like benign prostatic hyperplasia (BPH), neurogenic bladder syndrome, urinary incontinence like urge-,
10 stress-, or overflow incontinence, pelvic pain, and erectile dysfunction.

The human GSK3B is highly expressed in the following urological tissues: spinal cord (ventral horn), spinal cord (dorsal horn), fetal kidney, kidney, kidney, kidney tumor, renal epithelial cells, HEK 293 cells. The expression in the above mentioned tissues demonstrates that the human GSK3B or mRNA can be utilized to diagnose of urological disorders. Additionally the activity of
15 the human GSK3B can be modulated to treat urological disorders.

The human GSK3B is highly expressed in spinal cord tissues: spinal cord (ventral horn), spinal cord (dorsal horn). Expression in spinal cord tissues demonstrates that the human GSK3B or mRNA can be utilized to diagnose of incontinence as an urological disorder. The spinal cord tissues are involved in the neuronal regulation of the urological system. Additionally the activity of
20 the human GSK3B can be modulated to treat - but not limited to - incontinence.

Metabolic Disorders

Metabolic diseases are defined as conditions which result from an abnormality in any of the chemical or biochemical transformations and their regulating systems essential to producing energy, to regenerating cellular constituents, to eliminating unneeded products arising from these
25 processes, and to regulate and maintain homeostasis in a mammal regardless of whether acquired or the result of a genetic transformation. Depending on which metabolic pathway is involved, a single defective transformation or disturbance of its regulation may produce consequences that are narrow, involving a single body function, or broad, affecting many organs, organ-systems or the body as a whole. Diseases resulting from abnormalities related to the fine and coarse mechanisms
30 that affect each individual transformation, its rate and direction or the availability of substrates like amino acids, fatty acids, carbohydrates, minerals, cofactors, hormones, regardless whether they are

inborn or acquired, are well within the scope of the definition of a metabolic disease according to this application.

Metabolic diseases often are caused by single defects in particular biochemical pathways, defects that are due to the deficient activity of individual enzymes or molecular receptors leading to the regulation of such enzymes. Hence in a broader sense disturbances of the underlying genes, their products and their regulation lie well within the scope of this definition of a metabolic disease. For example, but not limited to, metabolic diseases may affect 1) biochemical processes and tissues ubiquitous all over the body, 2) the bone, 3) the nervous system, 4) the endocrine system, 5) the muscle including the heart, 6) the skin and nervous tissue, 7) the urogenital system, 8) the homeostasis of body systems like water and electrolytes. For example, but not limited to, metabolic diseases according to 1) comprise obesity, amyloidosis, disturbances of the amino acid metabolism like branched chain disease, hyperaminoacidemia, hyperaminoaciduria, disturbances of the metabolism of urea, hyperammonemia, mucopolysaccharidoses e.g. Maroteaux-Lamy syndrom, storage diseases like glycogen storage diseases and lipid storage diseases, glycogenosis diseases like Cori's disease, malabsorption diseases like intestinal carbohydrate malabsorption, oligosaccharidase deficiency like maltase-, lactase-, sucrase-insufficiency, disorders of the metabolism of fructose, disorders of the metabolism of galactose, galactosaemia, disturbances of carbohydrate utilization like diabetes, hypoglycemia, disturbances of pyruvate metabolism, hypolipidemia, hypolipoproteinemia, hyperlipidemia, hyperlipoproteinemia, carnitine or carnitine acyltransferase deficiency, disturbances of the porphyrin metabolism, porphyrias, disturbances of the purine metabolism, lysosomal diseases, metabolic diseases of nerves and nervous systems like gangliosidoses, sphingolipidoses, sulfatidoses, leucodystrophies, Lesch-Nyhan syndrome. For example, but not limited to, metabolic diseases according to 2) comprise osteoporosis, osteomalacia like osteoporosis, osteopenia, osteogenesis imperfecta, osteopetrosis, osteonecrosis, Paget's disease of bone, hypophosphatemia. For example, but not limited to, metabolic diseases according to 3) comprise cerebellar dysfunction, disturbances of brain metabolism like dementia, Alzheimer's disease, Huntington's chorea, Parkinson's disease, Pick's disease, toxic encephalopathy, demyelinating neuropathies like inflammatory neuropathy, Guillain-Barré syndrome. For example, but not limited to, metabolic diseases according to 4) comprise primary and secondary metabolic disorders associated with hormonal defects like any disorder stemming from either an hyperfunction or hypofunction of some hormone-secreting endocrine gland and any combination thereof. They comprise Sipple's syndrome, pituitary gland dysfunction and its effects on other endocrine glands, such as the thyroid, adrenals, ovaries, and testes, acromegaly, hyper- and hypothyroidism, euthyroid goiter, euthyroid sick syndrome, thyroiditis, and thyroid cancer, over- or underproduction of the adrenal steroid hormones, adrenogenital syndrome, Cushing's

syndrome, Addison's disease of the adrenal cortex, Addison's pernicious anemia, primary and secondary aldosteronism, diabetes insipidus, carcinoid syndrome, disturbances caused by the dysfunction of the parathyroid glands, pancreatic islet cell dysfunction, diabetes, disturbances of the endocrine system of the female like estrogen deficiency, resistant ovary syndrome. For example, but not limited to, metabolic diseases according to 5) comprise muscle weakness, myotonia, Duchenne's and other muscular dystrophies, dystrophia myotonica of Steinert, mitochondrial myopathies like disturbances of the catabolic metabolism in the muscle, carbohydrate and lipid storage myopathies, glycogenoses, myoglobinuria, malignant hyperthermia, polymyalgia rheumatica, dermatomyositis, primary myocardial disease, cardiomyopathy. For example, but not limited to, metabolic diseases according to 6) comprise disorders of the ectoderm, neurofibromatosis, scleroderma and polyarteritis, Louis-Bar syndrome, von Hippel-Lindau disease, Sturge-Weber syndrome, tuberous sclerosis, amyloidosis, porphyria. For example, but not limited to, metabolic diseases according to 7) comprise sexual dysfunction of the male and female. For example, but not limited to, metabolic diseases according to 8) comprise confused states and seizures due to inappropriate secretion of antidiuretic hormone from the pituitary gland, Liddle's syndrome, Bartter's syndrome, Fanconi's syndrome, renal electrolyte wasting, diabetes insipidus.

The human GSK3B is highly expressed in the following metabolic disease related tissues: thyroid tumor, cartilage and adipose. The expression in the above mentioned tissues demonstrates that the human GSK3B or mRNA can be utilized to diagnose of metabolic diseases. Additionally the activity of the human GSK3B can be modulated to treat metabolic diseases.

Applications

The present invention provides for both prophylactic and therapeutic methods for cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases.

The regulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of GSK3B. An agent that modulates activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of the polypeptide, a peptide, a peptidomimetic, or any small molecule. In one embodiment, the agent stimulates one or more of the biological activities of GSK3B. Examples of such stimulatory agents include the active GSK3B and nucleic acid molecules encoding a portion of GSK3B. In another embodiment, the agent inhibits one or more of the biological activities of GSK3B. Examples of such inhibitory agents include antisense nucleic acid molecules and antibodies. These regulatory methods can be performed in vitro (e.g., by culturing the cell with the agent) or, alternatively, in vivo (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating

an individual afflicted with a disease or disorder characterized by unwanted expression or activity of GSK3B or a protein in the GSK3B signaling pathway. In one embodiment, the method involves administering an agent like any agent identified or being identifiable by a screening assay as described herein, or combination of such agents that modulate say upregulate or downregulate the expression or activity of GSK3B or of any protein in the GSK3B signaling pathway. In another embodiment, the method involves administering a regulator of GSK3B as therapy to compensate for reduced or undesirably low expression or activity of GSK3B or a protein in the GSK3B signaling pathway.

Stimulation of activity or expression of GSK3B is desirable in situations in which enzymatic activity or expression is abnormally low and in which increased activity is likely to have a beneficial effect. Conversely, inhibition of enzymatic activity or expression of GSK3B is desirable in situations in which activity or expression of GSK3B is abnormally high and in which decreasing its activity is likely to have a beneficial effect.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

Pharmaceutical Compositions

This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

The nucleic acid molecules, polypeptides, and antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

The invention includes pharmaceutical compositions comprising a regulator of GSK3B expression or activity (and/or a regulator of the activity or expression of a protein in the GSK3B signaling pathway) as well as methods for preparing such compositions by combining one or more such

regulators and a pharmaceutically acceptable carrier. Also within the invention are pharmaceutical compositions comprising a regulator identified using the screening assays of the invention packaged with instructions for use. For regulators that are antagonists of GSK3B activity or which reduce GSK3B expression, the instructions would specify use of the pharmaceutical composition
5 for treatment of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases. For regulators that are agonists of GSK3B activity or increase GSK3B expression, the instructions would specify use of the pharmaceutical composition for treatment of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and
10 urological diseases.

An inhibitor of GSK3B may be produced using methods which are generally known in the art. In particular, purified GSK3B may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind GSK3B. Antibodies to GSK3B may also be generated using methods that are well known in the art. Such antibodies may include, but are
15 not limited to, polyclonal, monoclonal, chimeric, single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies like those which inhibit dimer formation are especially preferred for therapeutic use.

In another embodiment of the invention, the polynucleotides encoding GSK3B, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the
20 polynucleotide encoding GSK3B may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding GSK3B. Thus, complementary molecules or fragments may be used to modulate GSK3B activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can
25 be designed from various locations along the coding or control regions of sequences encoding GSK3B.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to
30 construct vectors which will express nucleic acid sequence complementary to the polynucleotides of the gene encoding GSK3B. These techniques are described, for example, in [Scott and Smith (1990)].

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

5 An additional embodiment of the invention relates to the administration of a pharmaceutical composition containing GSK3B in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of GSK3B, antibodies to GSK3B, and mimetics, agonists, antagonists, or inhibitors of GSK3B. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical
10 carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal
15 administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as
20 acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where
25 water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EMTM (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and
30 storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, a pharmaceutically acceptable polyol like glycerol, propylene glycol, liquid polyethylene glycol, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of

dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin. Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polypeptide or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for

transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

- 5 The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

10 In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with
15 monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to
20 physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations
25 inherent in the art of compounding such an active compound for the treatment of individuals.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration. For pharmaceutical compositions which include an antagonist of GSK3B activity, a compound which reduces expression of GSK3B, or a compound which reduces expression or activity of a protein in the GSK3B signaling pathway or any combination thereof,
30 the instructions for administration will specify use of the composition for cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases. For pharmaceutical compositions which include an agonist of GSK3B activity, a compound which increases expression of GSK3B, or a compound which increases expression or activity of a protein in the GSK3B signaling pathway or any

combination thereof, the instructions for administration will specify use of the composition for cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases.

Diagnostics

5 In another embodiment, antibodies which specifically bind GSK3B may be used for the diagnosis of disorders characterized by the expression of GSK3B, or in assays to monitor patients being treated with GSK3B or agonists, antagonists, and inhibitors of GSK3B. Antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for
10 to detect GSK3B in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent joining with a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring GSK3B, including ELISAs, RIAs, and FACS, are known in
15 the art and provide a basis for diagnosing altered or abnormal levels of GSK3B expression. Normal or standard values for GSK3B expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to GSK3B under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, preferably by photometric means. Quantities of GSK3B
20 expressed in subject samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding GSK3B may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect
25 and quantitate gene expression in biopsied tissues in which expression of GSK3B may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of GSK3B, and to monitor regulation of GSK3B levels during therapeutic intervention.

Polynucleotide sequences encoding GSK3B may be used for the diagnosis of cardiovascular
30 diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases associated with expression of GSK3B. The polynucleotide sequences encoding GSK3B may be used in Southern, Northern, or dot-blot analysis, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and

in microarrays utilizing fluids or tissues from patient biopsies to detect altered GSK3B expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding GSK3B may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding GSK3B may be labelled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding GSK3B in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

In order to provide a basis for the diagnosis of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases associated with expression of GSK3B, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding GSK3B, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

25 Determination of a Therapeutically Effective Dose

The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases GSK3B activity relative to GSK3B activity which occurs in the absence of the therapeutically effective dose. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

Therapeutic efficacy and toxicity, e.g., ED_{50} (the dose therapeutically effective in 50% of the population) and LD_{50} (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD_{50}/ED_{50} .

5 Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of
10 administration. The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction
15 sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts can vary from 0.1 micrograms to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and
20 methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc. If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either *ex vivo* or *in vivo* using well-
25 established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, "gene gun", and DEAE- or calcium phosphate-mediated transfection.

If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a
30 ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above. Preferably, a reagent reduces expression of GSK3B gene or the activity of GSK3B by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of GSK3B gene or the activity of
35 GSK3B can be assessed using methods well known in the art, such as hybridization of nucleotide

probes to GSK3B-specific mRNA, quantitative RT-PCR, immunologic detection of GSK3B, or measurement of GSK3B activity.

In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects. Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

Nucleic acid molecules of the invention are those nucleic acid molecules which are contained in a group of nucleic acid molecules consisting of (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1, (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1, (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code, wherein the polypeptide encoded by said nucleic acid molecule has GSK3B activity.

Polypeptides of the invention are those polypeptides which are contained in a group of polypeptides consisting of (i) polypeptides having the sequence of SEQ ID NO: 2, (ii) polypeptides comprising the sequence of SEQ ID NO: 2, (iii) polypeptides encoded by nucleic acid molecules of the invention and (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii), wherein said purified polypeptide has GSK3B activity.

An object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) contacting a test compound with a GSK3B polypeptide, (ii) detect binding of said test compound to said GSK3B polypeptide. E.g., compounds that bind to the GSK3B polypeptide are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) determining the activity of a GSK3B polypeptide at a certain concentration of a test compound or in the absence of said test compound, (ii) determining the activity of said polypeptide at a different concentration of said test compound. E.g., compounds that lead to a difference in the activity of the GSK3B polypeptide in (i) and (ii) are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) determining the activity of a GSK3B polypeptide at a certain concentration of a test compound, (ii) determining the activity of a GSK3B polypeptide at the presence of a compound known to be a regulator of a GSK3B polypeptide. E.g., compounds that show similar effects on the activity of the GSK3B polypeptide in (i) as compared to compounds used in (ii) are identified potential therapeutic agents for such a disease.

Other objects of the invention are methods of the above, wherein the step of contacting is in or at the surface of a cell.

Other objects of the invention are methods of the above, wherein the cell is in vitro.

Other objects of the invention are methods of the above, wherein the step of contacting is in a cell-free system.

Other objects of the invention are methods of the above, wherein the polypeptide is coupled to a detectable label.

Other objects of the invention are methods of the above, wherein the compound is coupled to a detectable label.

Other objects of the invention are methods of the above, wherein the test compound displaces a ligand which is first bound to the polypeptide.

Other objects of the invention are methods of the above, wherein the polypeptide is attached to a solid support.

Other objects of the invention are methods of the above, wherein the compound is attached to a solid support.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) contacting a test compound with a GSK3B polynucleotide, (ii) detect binding of said test compound to said GSK3B polynucleotide. Compounds that, e.g., bind to the GSK3B polynucleotide are potential therapeutic agents for the treatment of such diseases.

10 Another object of the invention is the method of the above, wherein the nucleic acid molecule is RNA.

Another object of the invention is a method of the above, wherein the contacting step is in or at the surface of a cell.

15 Another object of the invention is a method of the above, wherein the contacting step is in a cell-free system:

Another object of the invention is a method of the above, wherein the polynucleotide is coupled to a detectable label.

Another object of the invention is a method of the above, wherein the test compound is coupled to a detectable label.

20 Another object of the invention is a method of diagnosing a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) determining the amount of a GSK3B polynucleotide in a sample taken from said mammal, (ii) determining the amount of GSK3B polynucleotide in healthy and/or
25 diseased mammal. A disease is diagnosed, e.g., if there is a substantial similarity in the amount of GSK3B polynucleotide in said test mammal as compared to a diseased mammal.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological
30 diseases in a mammal comprising a therapeutic agent which binds to a GSK3B polypeptide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a GSK3B polypeptide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a GSK3B polypeptide, wherein said therapeutic agent is (i) a small molecule, (ii) an RNA molecule, (iii) an antisense oligonucleotide, (iv) a polypeptide, (v) an antibody, or (vi) a ribozyme.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a GSK3B polynucleotide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a GSK3B polypeptide.

Another object of the invention is the use of regulators of a GSK3B for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal.

Another object of the invention is a method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) identifying a regulator of GSK3B, (ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal; and (iii) combining of said regulator with an acceptable pharmaceutical carrier.

Another object of the invention is the use of a regulator of GSK3B for the regulation of GSK3B activity in a mammal having a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases.

- 5 The examples below are provided to illustrate the subject invention. These examples are provided by way of illustration and are not included for the purpose of limiting the invention.

Examples**Example 1: Search for homologous sequences in public sequence data bases**

The degree of homology can readily be calculated by known methods. Preferred methods to determine homology are designed to give the largest match between the sequences tested.

- 5 Methods to determine homology are codified in publicly available computer programs such as BestFit, BLASTP, BLASTN, and FASTA. The BLAST programs are publicly available from NCBI and other sources in the internet.

For GSK3B the following hits to known sequences were identified by using the BLAST algorithm [Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ; Nucleic Acids
10 Res 1997 Sep 1; 25(17): 3389-402] and the following set of parameters: matrix = BLOSUM62 and low complexity filter. The following databases were searched: NCBI (non-redundant database) and DERWENT patent database (Geneseq).

The following hits were found:

>gb|BC000251.1| Homo sapiens glycogen synthase kinase 3 beta, mRNA (cDNA cloneMGC:1736
15 IMAGE:3357620), complete.cds

Length = 1639, Score = 2711 bits (1410), Expect = 0.0, Identities = 1410/1410 (100%)

>ref|NM_002093.2| Homo sapiens glycogen synthase kinase 3 beta (GSK3B), mRNA

Length = 1639, Score = 2711 bits (1410), Expect = 0.0, Identities = 1410/1410 (100%)

>gb|AY335634.1| Synthetic construct Homo sapiens glycogen synthase kinase 3 beta(GSK3B)
20 mRNA, partial cds

Length = 1302, Score = 2500 bits (1300), Expect = 0.0, Identities = 1300/1300 (100%)

>gb|AR270851.1| Sequence 1414 from patent US 6500938

Length = 1389, Score = 1790 bits (931), Expect = 0.0, Identities = 931/931 (100%)

>gb|AR262205.1| Sequence 3 from patent US 6323029

25 Length = 1389, Score = 1790 bits (931), Expect = 0.0, Identities = 931/931 (100%)

>emb|AX821914.1| Sequence 42 from Patent WO03068961

Length = 1231, Score = 1790 bits (931), Expect = 0.0, Identities = 931/931 (100%)

>emb|AX701653.1| Sequence 3 from Patent WO03000882

Length = 1389, Score = 1790 bits (931), Expect = 0.0, Identities = 931/931 (100%)

>emb|AX777402.1| Sequence 256 from Patent WO03040301

5 Length = 1389, Score = 1790 bits (931), Expect = 0.0, Identities = 931/931 (100%)

>gb|BC012760.2| Homo sapiens glycogen synthase kinase 3 beta, mRNA (cDNA cloneMGC:16182 IMAGE:3637163), complete cds

Length = 2374, Score = 1790 bits (931), Expect = 0.0, Identities = 931/931 (100%)

>gb|L33801.1|HUMGLSYKIN Human protein kinase mRNA, complete cds

10 Length = 1389, Score = 1790 bits (931), Expect = 0.0, Identities = 931/931 (100%)

>gb|AR059073.1|AR059073 Sequence 1 from patent US 5837853

Length = 2088, Score = 1779 bits (925), Expect = 0.0, Identities = 929/931 (99%)

>gb|AR097210.1|AR097210 Sequence 1 from patent US 6071694

Length = 2088, Score = 1779 bits (925), Expect = 0.0, Identities = 929/931 (99%)

15 >dbj|E08052.1| cDNA encoding human tauproteinkinase-1

Length = 2088, Score = 1779 bits (925), Expect = 0.0, Identities = 929/931 (99%)

>emb|AX701656.1| Sequence 6 from Patent WO03000882

Length = 1263, Score = 1750 bits (910), Expect = 0.0, Identities = 910/910 (100%)

>ref|NM_019827.2| Mus musculus glycogen synthase kinase 3 beta (GSK3B), mRNA

20 Length = 2841, Score = 1392 bits (724), Expect = 0.0, Identities = 850/913 (93%)

>gb|BC060743.1| Mus musculus glycogen synthase kinase 3 beta, mRNA (cDNA cloneMGC:68385 IMAGE:4022374), complete cds

Length = 2841, Score = 1392 bits (724), Expect = 0.0, Identities = 850/913 (93%)

>gb|BC006936.1| Mus musculus glycogen synthase kinase 3 beta, mRNA (cDNA clone MGC:6814 IMAGE:2648507), complete cds

Length = 1503, Score = 1392 bits (724), Expect = 0.0, Identities = 850/913 (93%)

>gb|AF156099.2|AF156099 Mus musculus glycogen synthase kinase 3 beta mRNA, complete cds

5 Length = 1535, Score = 1392 bits (724), Expect = 0.0, Identities = 850/913 (93%)

>gb|AR059074.1|AR059074 Sequence 2 from patent US 5837853

Length = 1972, Score = 1363 bits (709), Expect = 0.0, Identities = 845/913 (92%)

>gb|AR097211.1|AR097211 Sequence 2 from patent US 6071694

Length = 1972, Score = 1363 bits (709), Expect = 0.0, Identities = 845/913 (92%)

10 >dbj|BD181611.1| Method of the phosphorylation of tau protein

Length = 1972, Score = 1363 bits (709), Expect = 0.0, Identities = 845/913 (92%)

>dbj|E08007.1| DNA encoding tau protein kinase I

Length = 1972, Score = 1363 bits (709), Expect = 0.0, Identities = 845/913 (92%)

>emb|X73653.1|RNTAU R. norvegicus mRNA for tau protein kinase I

15 Length = 1525, Score = 1363 bits (709), Expect = 0.0, Identities = 845/913 (92%)

>ref|NM_032080.1| Rattus norvegicus glycogen synthase kinase 3 beta (GSK3B), mRNA

Length = 1525, Score = 1363 bits (709), Expect = 0.0, Identities = 845/913 (92%)

>emb|X53428.1|RNGSK3B Rat mRNA for glycogen synthase kinase 3 beta (EC 2.7.1.37)

Length = 1474, Score = 1352 bits (703), Expect = 0.0, Identities = 843/913 (92%)

20 >gb|AR059082.1|AR059082 Sequence 13 from patent US 5837853

Length = 479, Score = 892 bits (464), Expect = 0.0, Identities = 474/479 (98%)

>gb|AR097219.1|AR097219 Sequence 13 from patent US 6071694

Length = 479, Score = 892 bits (464), Expect = 0.0, Identities = 474/479 (98%)

>dbj|E08056.1| cDNA encoding part of human tauproteinkinase-1

Length = 482, Score = 888 bits (462), Expect = 0.0, Identities = 477/482 (98%), Gaps = 3/482 (0%).

Example 2: Expression profiling

- 5 Total cellular RNA was isolated from cells by one of two standard methods: 1) guanidine isothiocyanate/Cesium chloride density gradient centrifugation [Kellogg, (1990)]; or with the Tri-Reagent protocol according to the manufacturer's specifications (Molecular Research Center, Inc., Cincinnati, Ohio). Total RNA prepared by the Tri-reagent protocol was treated with DNase I to remove genomic DNA contamination.
- 10 For relative quantitation of the mRNA distribution of GSK3B, total RNA from each cell or tissue source was first reverse transcribed. 85 µg of total RNA was reverse transcribed using 1 µmole random hexamer primers, 0.5 mM each of dATP, dCTP, dGTP and dTTP (Qiagen, Hilden, Germany), 3000 U RnaseQut (Invitrogen, Groningen, Netherlands) in a final volume of 680 µl. The first strand synthesis buffer and Omniscript reverse transcriptase (2 u/µl) were from (Qiagen,
- 15 Hilden, Germany). The reaction was incubated at 37°C for 90 minutes and cooled on ice. The volume was adjusted to 6800 µl with water, yielding a final concentration of 12.5 ng/µl of starting RNA.

- For relative quantitation of the distribution of GSK3B mRNA in cells and tissues the Perkin Elmer ABI Prism RTM. 7700 Sequence Detection system or Biorad iCycler was used according to the
- 20 manufacturer's specifications and protocols. PCR reactions were set up to quantitate GSK3B and the housekeeping genes HPRT (hypoxanthine phosphoribosyltransferase), GAPDH (glyceraldehyde-3-phosphate dehydrogenase), β-actin, and others. Forward and reverse primers and probes for GSK3B were designed using the Perkin Elmer ABI Primer Express™ software and were synthesized by TibMolBiol (Berlin, Germany). The GSK3B forward primer sequence was:
- 25 Primer1 (SEQ ID NO: 3). The GSK3B reverse primer sequence was Primer2 (SEQ ID NO: 4). Probe1 (SEQ ID NO: 5), labelled with FAM (carboxyfluorescein succinimidyl ester) as the reporter dye and TAMRA (carboxytetramethylrhodamine) as the quencher, is used as a probe for GSK3B. The following reagents were prepared in a total of 25 µl : 1x TaqMan buffer A, 5.5 mM MgCl₂, 200 nM of dATP, dCTP, dGTP, and dUTP, 0.025 U/µl AmpliTaq Gold™, 0.01 U/ µl
- 30 AmpErase and Probe1 (SEQ ID NO: 5), GSK3B forward and reverse primers each at 200 nM, 200 nM GSK3B FAM/TAMRA-labelled probe, and 5 µl of template cDNA. Thermal cycling parameters were 2 min at 50°C, followed by 10 min at 95°C, followed by 40 cycles of melting at 95°C for 15 sec and annealing/extending at 60°C for 1 min.

Calculation of corrected CT values

The CT (threshold cycle) value is calculated as described in the "Quantitative determination of nucleic acids" section. The CF-value (factor for threshold cycle correction) is calculated as follows :

- 5 1. PCR reactions were set up to quantitate the housekeeping genes (HKG) for each cDNA sample.
2. CT_{HKG} -values (threshold cycle for housekeeping gene) were calculated as described in the "Quantitative determination of nucleic acids" section.
3. CT_{HKG} -mean values (CT mean value of all HKG tested on one cDNAs) of all HKG for
10 each cDNA are calculated (n = number of HKG):

$$CT_{HKG-n}\text{-mean value} = (CT_{HKG1}\text{-value} + CT_{HKG2}\text{-value} + \dots + CT_{HKG-n}\text{-value}) / n$$
4. CT_{panel} mean value (CT mean value of all HKG in all tested cDNAs) =

$$(CT_{HKG1}\text{-mean value} + CT_{HKG2}\text{-mean value} + \dots + CT_{HKG-y}\text{-mean value}) / y$$
 (y = number of cDNAs)
- 15 5. CF_{cDNA-n} (correction factor for cDNA n) = $CT_{\text{panel}}\text{-mean value} - CT_{HKG-n}\text{-mean value}$
6. CT_{cDNA-n} (CT value of the tested gene for the cDNA n) + CF_{cDNA-n} (correction factor for cDNA n) = $CT_{\text{cor-cDNA-n}}$ (corrected CT value for a gene on cDNA n)

Calculation of relative expression

Definition : highest $CT_{\text{cor-cDNA-n}} \neq 40$ is defined as $CT_{\text{cor-cDNA}} [\text{high}]$

20 $\text{Relative Expression} = 2^{(CT_{\text{cor-cDNA}}[\text{high}] - CT_{\text{cor-cDNA-n}})}$

Tissues

The expression of GSK3B was investigated in the tissues in table 1.

Expression profile

The results of the the mRNA-quantification (expression profiling) is shown in Table 1.

Table 1: Relative expression of GSK3B in various human tissues.

T-cells peripheral blood CD4+	101
T-cells peripheral blood CD4+	38
T-cells peripheral blood CD4+ D117 II virus infected	36
T-cells peripheral blood CD4+ D34 virus infected	41
monocytes	171
monocytes HIV-1 infected	190
fetal heart	43
heart	16
heart	30
heart	15
heart	208
heart myocardial infarction	315
heart myocardial infarction	198
heart myocardial infarction	199
pericardium	193
heart atrium (right)	171
heart atrium (right)	161
heart atrium (left)	242
heart atrium (left)	164
heart ventricle (left)	37
heart ventricle (left)	174
heart ventricle (right)	56
heart ventricle (right)	241
heart apex	126

Purkinje fibers	185
interventricular septum	242
fetal aorta	22
aorta	17
aorta	16
aorta	10
arcus aorta	24
aorta valve	519
artery	9
coronary artery	16
coronary artery	14
coronary artery	12
pulmonary artery	10
carotid artery	9
mesenteric artery	13
arteria radialis	12
vein	13
pulmonic valve	45
vein (saphena magna)	16
(caval) vein	9
coronary artery endothel cells	142
coronary artery smooth muscle primary cells	66
aortic smooth muscle cells	120
pulmonary artery smooth muscle cells	111
aortic endothel cells	207
HUVEC cells	114
pulmonary artery endothel cells	163
iliac artery endothel cells	169

skin	168
adrenal gland	140
thyroid	131
thyroid tumor	197
pancreas	25
pancreas liver cirrhosis	19
esophagus	15
esophagus tumor	182
stomach	121
stomach tumor	103
colon	87
colon tumor	91
small intestine	82
ileum	117
ileum tumor	64
ileum chronic inflammation	0
rectum	117
rectum tumor	290
fetal liver	70
liver	89
liver	5
liver	3
liver liver cirrhosis	64
liver lupus disease	137
liver tumor	174

HEP G2 cells	234
leukocytes (peripheral blood)	68
Jurkat (T-cells)	133
Raji (B-cells)	115
bone marrow	41
erythrocytes	3
lymphnode	11
thymus	104
thrombocytes	9
bone marrow stromal cells	65
bone marrow CD71+ cells	2
bone marrow CD33+ cells	10
bone marrow CD34+ cells	19
bone marrow CD15+ cells	3
cord blood CD71+ cells	1
cord blood CD34+ cells	70
neutrophils cord blood	111
T-cells peripheral blood CD8+	109
monocytes peripheral blood CD14+	145
B-cells peripheral blood CD19+	60
neutrophils peripheral blood	360
spleen	128
spleen liver cirrhosis	54
skeletal muscle	122
cartilage	181
bone connective tissue	221

adipose	48
adipose	143
adipose	236
fetal adipose	383
adipose (subcutaneous) BMI 21.74	7
adipose (subcutaneous) BMI 35.04	2
brain	232
cerebellum	81
cerebral cortex	335
frontal lobe	474
occipital lobe	422
parietal lobe	530
temporal lobe	671
substantia nigra	50
caudatum	393
corpus callosum	202
nucleus accumbens	580
putamen	443
hippocampus	393
thalamus	159
posteroventral thalamus	474
dorsalmedial thalamus	449
hypothalamus	355
dorsal root ganglia	14
spinal cord	151
spinal cord (ventral horn)	286
spinal cord (dorsal horn)	350

glial tumor H4 cells	110
astrocytes	236
retina	36
fetal lung	138
fetal lung fibroblast IMR-90 cells	175
fetal lung fibroblast MRC-5 cells	167
lung	14
lung	22
lung	4
lung right upper lobe	45
lung right mid lobe	63
lung right lower lobe	55
lung lupus disease	47
lung tumor	159
lung COPD	9
trachea	99
primary bronchia	139
secondary bronchia	96
bronchial epithelial cells	333
bronchial smooth muscle cells	153
small airway epithelial cells	534
cervix	26
testis	347
HeLa cells (cervix tumor)	105
placenta	133
uterus	134

uterus tumor	132
ovary	120
ovary tumor	258
breast	111
breast tumor	82
mammary gland	120
prostate	138
prostate	265
prostate	114
prostate BPH	10
prostate tumor	184
bladder	72
bladder	184
bladder	147
ureter	28
penis	6
corpus cavernosum	18
fetal kidney	232
kidney	114
kidney	17
kidney	150
kidney tumor	136
renal epithelial cells	175
HEK 293 cells	162

Example 3: Antisense Analysis

Knowledge of the correct, complete cDNA sequence coding for GSK3B enables its use as a tool for antisense technology in the investigation of gene function. Oligonucleotides, cDNA or genomic fragments comprising the antisense strand of a polynucleotide coding for GSK3B are used either in vitro or in vivo to inhibit translation of the mRNA. Such technology is now well known in the art, and antisense molecules can be designed at various locations along the nucleotide sequences. By treatment of cells or whole test animals with such antisense sequences, the gene of interest is effectively turned off. Frequently, the function of the gene is ascertained by observing behavior at the intracellular, cellular, tissue or organismal level (e.g., lethality, loss of differentiated function, changes in morphology, etc.).

In addition to using sequences constructed to interrupt transcription of a particular open reading frame, modifications of gene expression is obtained by designing antisense sequences to intron regions, promoter/enhancer elements, or even to trans-acting regulatory genes.

Example 4: Expression of GSK3B

Expression of GSK3B is accomplished by subcloning the cDNAs into appropriate expression vectors and transfecting the vectors into expression hosts such as, e.g., *E. coli*. In a particular case, the vector is engineered such that it contains a promoter for β -galactosidase, upstream of the cloning site, followed by sequence containing the amino-terminal Methionine and the subsequent seven residues of β -galactosidase. Immediately following these eight residues is an engineered bacteriophage promoter useful for artificial priming and transcription and for providing a number of unique endonuclease restriction sites for cloning.

Induction of the isolated, transfected bacterial strain with Isopropyl- β -D-thiogalactopyranoside (IPTG) using standard methods produces a fusion protein corresponding to the first seven residues of β -galactosidase, about 15 residues of "linker", and the peptide encoded within the cDNA. Since cDNA clone inserts are generated by an essentially random process, there is probability of 33% that the included cDNA will lie in the correct reading frame for proper translation. If the cDNA is not in the proper reading frame, it is obtained by deletion or insertion of the appropriate number of bases using well known methods including in vitro mutagenesis, digestion with exonuclease III or mung bean nuclease, or the inclusion of an oligonucleotide linker of appropriate length.

The GSK3B cDNA is shuttled into other vectors known to be useful for expression of proteins in specific hosts. Oligonucleotide primers containing cloning sites as well as a segment of DNA (about 25 bases) sufficient to hybridize to stretches at both ends of the target cDNA is synthesized

chemically by standard methods. These primers are then used to amplify the desired gene segment by PCR. The resulting gene segment is digested with appropriate restriction enzymes under standard conditions and isolated by gel electrophoresis. Alternately, similar gene segments are produced by digestion of the cDNA with appropriate restriction enzymes. Using appropriate
5 primers, segments of coding sequence from more than one gene are ligated together and cloned in appropriate vectors. It is possible to optimize expression by construction of such chimeric sequences.

Suitable expression hosts for such chimeric molecules include, but are not limited to, mammalian cells such as Chinese Hamster Ovary (CHO) and human 293 cells; insect cells such as Sf9 cells,
10 yeast cells such as *Saccharomyces cerevisiae* and bacterial cells such as *E. coli*. For each of these cell systems, a useful expression vector also includes an origin of replication to allow propagation in bacteria, and a selectable marker such as the β -lactamase antibiotic resistance gene to allow plasmid selection in bacteria. In addition, the vector may include a second selectable marker such as the neomycin phosphotransferase gene to allow selection in transfected eukaryotic host cells.
15 Vectors for use in eukaryotic expression hosts require RNA processing elements such as 3' polyadenylation sequences if such are not part of the cDNA of interest.

Additionally, the vector contains promoters or enhancers which increase gene expression. Such promoters are host specific and include MMTV, SV40, and metallothionine promoters for CHO cells; trp, lac, tac and T7 promoters for bacterial hosts; and alpha factor, alcohol oxidase and PGH
20 promoters for yeast. Transcription enhancers, such as the rous sarcoma virus enhancer, are used in mammalian host cells. Once homogeneous cultures of recombinant cells are obtained through standard culture methods, large quantities of recombinantly produced GSK3B are recovered from the conditioned medium and analyzed using chromatographic methods known in the art. For example, GSK3B can be cloned into the expression vector pcDNA3, as exemplified herein. This
25 product can be used to transform, for example, HEK293 or COS by methodology standard in the art. Specifically, for example, using Lipofectamine (Gibco BRL catalog no. 18324-020) mediated gene transfer.

Example 5: Isolation of Recombinant GSK3B

GSK3B is expressed as a chimeric protein with one or more additional polypeptide domains added
30 to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals [Appa Rao, 1997] and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Washington). The inclusion of a cleavable linker

sequence such as Factor Xa or enterokinase (Invitrogen, Groningen, The Netherlands) between the purification domain and the GSK3B sequence is useful to facilitate expression of GSK3B.

The following example provides a method for purifying GSK3B.

5 GSK3B is generated using the baculovirus expression system BAC-TO-BAC (GIBCO BRL) based on *Autographa californica* nuclear polyhedrosis virus (AcNPV) infection of *Spodoptera frugiperda* insect cells (Sf9 cells).

10 cDNA encoding kinases cloned into either the donor plasmid pFASTBAC1 or pFASTBAC-HT which contain a mini-Tn7 transposition element. The recombinant plasmid is transformed into DH10BAC competent cells which contain the parent bacmid bMON14272 (AcNPV infectious DNA) and a helper plasmid. The mini-Tn7 element on the pFASTBAC donor can transpose to the attTn7 attachment site on the bacmid thus introducing the kinase gene into the viral genome. Colonies containing recombinant bacmids are identified by disruption of the *lacZ* gene. The kinase/bacmid construct can then be isolated and infected into insect cells (Sf9 cells) resulting in the production of infectious recombinant baculovirus particles and expression of either unfused
15 recombinant enzyme (pFastbac1) or GSK3B-His fusion protein (pFastbacHT).

Cells are harvested and extracts prepared 24, 48 and 72 hours after transfection. Expression of GSK3B is confirmed by coomassie staining after sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) and western blotting onto a PVDF membrane of an unstained SDS-PAGE. The kinase-His fusion protein is detected due to the interaction between the Ni-NTA HRP
20 conjugate and the His-tag which is fused to GSK3B.

Example 6: Production of GSK3B Specific Antibodies

Two approaches are utilized to raise antibodies to GSK3B, and each approach is useful for generating either polyclonal or monoclonal antibodies. In one approach, denatured protein from reverse phase HPLC separation is obtained in quantities up to 75 mg. This denatured protein is
25 used to immunize mice or rabbits using standard protocols; about 100 µg are adequate for immunization of a mouse, while up to 1 mg might be used to immunize a rabbit. For identifying mouse hybridomas, the denatured protein is radioiodinated and used to screen potential murine B-cell hybridomas for those which produce antibody. This procedure requires only small quantities of protein, such that 20 mg is sufficient for labeling and screening of several thousand clones.

30 In the second approach, the amino acid sequence of an appropriate GSK3B domain, as deduced from translation of the cDNA, is analyzed to determine regions of high antigenicity. Oligopeptides comprising appropriate hydrophilic regions are synthesized and used in suitable immunization

protocols to raise antibodies. The optimal amino acid sequences for immunization are usually at the C-terminus, the N-terminus and those intervening, hydrophilic regions of the polypeptide which are likely to be exposed to the external environment when the protein is in its natural conformation.

- 5 Typically, selected peptides, about 15 residues in length, are synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to keyhole limpet hemocyanin (KLH; Sigma, St. Louis, MO) by reaction with M-maleimidobenzoyl-N-hydroxy-succinimide ester, MBS. If necessary, a cysteine is introduced at the N-terminus of the peptide to permit coupling to KLH. Rabbits are immunized with the peptide-KLH complex in complete
10 Freund's adjuvant. The resulting antisera are tested for antipeptide activity by binding the peptide to plastic, blocking with 1% bovine serum albumin, reacting with antisera, washing and reacting with labeled (radioactive or fluorescent), affinity purified, specific goat anti-rabbit IgG.

- Hybridomas are prepared and screened using standard techniques. Hybridomas of interest are detected by screening with labeled GSK3B to identify those fusions producing the monoclonal
15 antibody with the desired specificity. In a typical protocol, wells of plates (FAST; Becton-Dickinson, Palo Alto, CA) are coated during incubation with affinity purified, specific rabbit anti-mouse (or suitable antispecies 1 g) antibodies at 10 mg/ml. The coated wells are blocked with 1% bovine serum albumin, (BSA), washed and incubated with supernatants from hybridomas. After washing the wells are incubated with labeled GSK3B at 1 mg/ml. Supernatants with specific
20 antibodies bind more labeled GSK3B than is detectable in the background. Then clones producing specific antibodies are expanded and subjected to two cycles of cloning at limiting dilution. Cloned hybridomas are injected into pristane-treated mice to produce ascites, and monoclonal antibody is purified from mouse ascitic fluid by affinity chromatography on Protein A. Monoclonal antibodies with affinities of at least
25 10^8 M^{-1} , preferably 10^9 to 10^{10} M^{-1} or stronger, are typically made by standard procedures.

Example 7: Diagnostic Test Using GSK3B Specific Antibodies

Particular GSK3B antibodies are useful for investigating signal transduction and the diagnosis of infectious or hereditary conditions which are characterized by differences in the amount or distribution of GSK3B or downstream products of an active signaling cascade.

- 30 Diagnostic tests for GSK3B include methods utilizing antibody and a label to detect GSK3B in human body fluids, membranes, cells, tissues or extracts of such. The polypeptides and antibodies of the present invention are used with or without modification. Frequently, the polypeptides and

antibodies are labeled by joining them, either covalently or noncovalently, with a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and have been reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemi-
5 luminescent agents, chromogenic agents, magnetic particles and the like.

A variety of protocols for measuring soluble or membrane-bound GSK3B, using either polyclonal or monoclonal antibodies specific for the protein, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site monoclonal-based immunoassay utilizing monoclonal antibodies
10 reactive to two non-interfering epitopes on GSK3B is preferred, but a competitive binding assay may be employed.

Example 8: Purification of Native GSK3B Using Specific Antibodies

Native or recombinant GSK3B is purified by immunoaffinity chromatography using antibodies specific for GSK3B. In general, an immunoaffinity column is constructed by covalently coupling
15 the anti-TRH antibody to an activated chromatographic resin.

Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified
20 immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated Sepharose (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such immunoaffinity columns are utilized in the purification of GSK3B by preparing a fraction from cells containing GSK3B in a soluble form. This preparation is derived by solubilization of
25 whole cells or of a subcellular fraction obtained via differential centrifugation (with or without addition of detergent) or by other methods well known in the art. Alternatively, soluble GSK3B containing a signal sequence is secreted in useful quantity into the medium in which the cells are grown:

A soluble GSK3B-containing preparation is passed over the immunoaffinity column, and the
30 column is washed under conditions that allow the preferential absorbance of GSK3B (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions

that disrupt antibody/protein binding (e.g., a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and GSK3B is collected.

Example 9: Drug Screening

This invention is particularly useful for screening therapeutic compounds by using GSK3B or
5 fragments thereof in any of a variety of drug screening techniques.

The following example provides a system for drug screening measuring the kinase activity.

The recombinant kinase-His fusion protein can be purified from the crude lysate by metal-affinity chromatography using Ni-NTA agarose. This allows the specific retention of the recombinant material (since this is fused to the His-tag) whilst the endogenous insect proteins are washed off.

10 The recombinant material is then eluted by competition with imidazol.

The activity of GSK3B molecules of the present invention can be measured using a variety of assays that measure GSK3B activity. For example, GSK3B enzyme activity can be assessed by a standard *in vitro* kinase assay.

The kinase activity of the kinase can be detected, for example, by adding ATP having radioactively
15 labeled phosphate to the reaction system containing the protein of the present invention and the substrate and measuring the radioactivity of the phosphate attached to the substrate.

Example 10: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact, agonists, antagonists, or inhibitors. Any
20 of these examples are used to fashion drugs which are more active or stable forms of the polypeptide or which enhance or interfere with the function of a polypeptide *in vivo*.

In one approach, the three-dimensional structure of a protein of interest, or of a protein-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the polypeptide must be
25 ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of a polypeptide is gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design efficient inhibitors. Useful examples of rational drug design include molecules which have improved activity or stability or which act as inhibitors, agonists, or antagonists of native peptides.

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design is based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original receptor. The anti-id is then used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides then act as the pharmacore.

By virtue of the present invention, sufficient amount of polypeptide are made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the GSK3B amino acid sequence provided herein provides guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

Example 11: Identification of Other Members of the Signal Transduction Complex

Labeled GSK3B is useful as a reagent for the purification of molecules with which it interacts. In one embodiment of affinity purification, GSK3B is covalently coupled to a chromatography column. Cell-free extract derived from synovial cells or putative target cells is passed over the column, and molecules with appropriate affinity bind to GSK3B. GSK3B-complex is recovered from the column, and the GSK3B-binding ligand disassociated and subjected to N-terminal protein sequencing. The amino acid sequence information is then used to identify the captured molecule or to design degenerate oligonucleotide probes for cloning the relevant gene from an appropriate cDNA library.

In an alternate method, antibodies are raised against GSK3B, specifically monoclonal antibodies. The monoclonal antibodies are screened to identify those which inhibit the binding of labeled GSK3B. These monoclonal antibodies are then used therapeutically.

Example 12: Use and Administration of Antibodies, Inhibitors, or Antagonists

Antibodies, inhibitors, or antagonists of GSK3B or other treatments and compounds that are limiters of signal transduction (LSTs), provide different effects when administered therapeutically. LSTs are formulated in a nontoxic, inert, pharmaceutically acceptable aqueous carrier medium preferably at a pH of about 5 to 8, more preferably 6 to 8, although pH may vary according to the characteristics of the antibody, inhibitor, or antagonist being formulated and the condition to be treated. Characteristics of LSTs include solubility of the molecule, its half-life and antigenicity/-immunogenicity. These and other characteristics aid in defining an effective carrier. Native

human proteins are preferred as LSTs, but organic or synthetic molecules resulting from drug screens are equally effective in particular situations.

5 LSTs are delivered by known routes of administration including but not limited to topical creams and gels; transmucosal spray and aerosol; transdermal patch and bandage; injectable, intravenous and lavage formulations; and orally administered liquids and pills particularly formulated to resist stomach acid and enzymes. The particular formulation, exact dosage, and route of administration is determined by the attending physician and varies according to each specific situation.

10 Such determinations are made by considering multiple variables such as the condition to be treated, the LST to be administered, and the pharmacokinetic profile of a particular LST. Additional factors which are taken into account include severity of the disease state, patient's age, weight, gender and diet, time and frequency of LST administration, possible combination with other drugs, reaction sensitivities, and tolerance/response to therapy. Long acting LST formulations might be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular LST.

15 Normal dosage amounts vary from 0.1 to 10^5 μ g, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. Those skilled in the art employ different formulations for different LSTs. Administration to cells such as nerve cells necessitates delivery in a manner different from that to other cells such as vascular endothelial
20 cells.

It is contemplated that abnormal signal transduction, trauma, or diseases which trigger GSK3B activity are treatable with LSTs. These conditions or diseases are specifically diagnosed by the tests discussed above, and such testing should be performed in suspected cases of viral, bacterial or fungal infections, allergic responses, mechanical injury associated with trauma, hereditary
25 diseases, lymphoma or carcinoma, or other conditions which activate the genes of lymphoid or neuronal tissues.

Example 13: Production of Non-human Transgenic Animals

Animal model systems which elucidate the physiological and behavioral roles of the GSK3B are produced by creating nonhuman transgenic animals in which the activity of the GSK3B is either
30 increased or decreased, or the amino acid sequence of the expressed GSK3B is altered, by a variety of techniques. Examples of these techniques include, but are not limited to: 1) Insertion of normal or mutant versions of DNA encoding a GSK3B, by microinjection, electroporation, retroviral

transfection or other means well known to those skilled in the art, into appropriately fertilized embryos in order to produce a transgenic animal or 2) homologous recombination of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these GSK3B sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and hence is useful for producing an animal that cannot express native GSK3Bs but does express, for example, an inserted mutant GSK3B, which has replaced the native GSK3B in the animal's genome by recombination, resulting in underexpression of the kinase. Microinjection adds genes to the genome, but does not remove them, and the technique is useful for producing an animal which expresses its own and added GSK3B, resulting in overexpression of the GSK3B.

One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as cesiumchloride M2 medium. DNA or cDNA encoding GSK3B is purified from a vector by methods well known to the one skilled in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the transgene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the transgene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a piper puller) and the egg to be injected is put in a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse which is a mouse stimulated by the appropriate hormones in order to maintain false pregnancy, where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only method for inserting DNA into the egg but is used here only for exemplary purposes.

25 References

- U.S. 4,522,811
- U.S. 5,283,317
- U.S. 5,565,332
- U.S. 5,723,323
- 30 U.S. 5,747,334
- U.S. 5,783,384
- U.S. 5,837,853

- U.S. 6,071,694
- U.S. 6,323,029
- U.S. 6,500,938
- WO 84/03564
- 5 WO 93/03151
- WO 94/13804
- WO 03/000882
- WO 03/040301
- WO 03/068961
- 10 Altschul SF et al; Nucleic Acids Res 1997 Sep 1; 25(17): 3389-402
- Appa Rao et al., 1997, Protein Expr Purif Nov, 11(2): 201-8
- Avalle et al., Ann. N Y Acad.Sci. 864:118 (1998)).
- Barnes, 2000, Chest, 117:10S14S
- Botstein et al., 1980, Am J Hum Genet. 32: 314-31
- 15 Colbere-Garapin *et al.*, 1981, *J. Mol. Biol.* 150, 1-14
- Cunningham and Wells, *J. Mol. Biol.* 234:554 (1993).
- Egan, S. E. and Weinberg, R. A. (1993) *Nature* 365:781-783
- Engelhard *et al.*, 1994, *Proc. Nat. Acad. Sci.* 91, 3224-3227
- Frame et al. (2001), *Molec. Cell* 7: 1321-1327
- 20 Friboulet et al., *Appl. Biochem. Biotechnol.* 47:229 (1994)
- Gao, G. et al. (1996) *J. Biol Chem.* 15:8675-81
- Gergen and Weiss , 1992, *Am Rev Respir Dis* 146:823-824
- Gibson et al., 1996, *Genome Research* 6: 995-1001
- Hardie, G. and Hanks, S. (1995) *The Protein Kinase Facts Books*, Vol I:7-20 Academic Press, San Diego, Calif.
- 25 Haribabu, B. et al. (1995) *EMBO Journal* 14:3679-86
- Harlow, *Antibodies*, Cold Spring Harbor Press, (1989)
- Haseloff *et al.*, 1988 , *Nature* 334, 585-591
- Heid et al., 1996, *Genome Research* 6: 986-994

- Holland et al., 1991, PNAS 88: 7276-7280
- Isselbacher, K. J. et al. (1994) Harrison's Principles of Internal Medicine, McGraw-Hill, New York, N.Y., pp. 416-431, 1887
- Jeffreys et al., 1985, Nature 316: 76-9
- 5 Johnson et al., 1989, *Endoc. Rev.* 10, 317-331
- Joron et al., Ann. N Y Acad. Sci. 672:216 (1992)
- Karlsson, Immunol. Methods 145:229 (1991)
- Kellogg et al., 1990, Anal. Biochem. 189:202-208
- Lam , 1997, Anticancer Drug Res. 12(3):145-67
- 10 Li, B. et al. (1996) J. Biol. Chem. 271:19402-8
- Livak et al., 1995 , PCR Methods and Applications 357-362
- Logan, Shenk, 1984, *Proc. Natl. Acad. Sci.* 81, 3655-3659
- Lowy et al., 1980, *Cell* 22, 817-23
- Maddox et al., 1983, *J. Exp. Med.* 158, 1211-1216
- 15 McConnell et al., 1992 , *Science* 257, 1906-1912
- Monfardini et al., Proc. Assoc. Am. Physicians 108:420 (1996)
- Nicholls et al., 1993, *J. Immunol. Meth.* 165, 81-91
- Piatak et al., 1993, BioTechniques 14:70-81
- Piatak et al., 1993, Science 259:1749-1754
- 20 Plyte et al., (1992), *Biochim. Biophys. Acta* 1114: 147-162
- Porath et al., 1992, *Prot. Exp. Purif.* 3, 263-281
- Roberge et al., 1995, *Science* 269, 202-204
- Scott and Smith (1990) Science 249:386-390
- Sjolander, Urbaniczky, 1991, *Anal. Chem.* 63, 2338-2345
- 25 Stambolic and Woodgett, (1994), *Biochem. J.* 303: 701-704
- Szabo et al., 1995, *Curr. Opin. Struct. Biol.* 5, 699-705
- Thomas, 1980, Proc. Nat. Acad. Sci., 77:5201-5205
- Uhlmann et al., 1987, *Tetrahedron. Lett.* 215, 3539-3542
- Weber et al., 1990, Genomics 7: 524-30

Wigler *et al.*, 1977, *Cell* 11, 223-32

Wigler *et al.*, 1980, *Proc. Natl. Acad. Sci.* 77, 3567-70

Woodgett (1990), *EMBO J.* 9: 2431-2438

26. Feb. 2004

Claims

1. A method of screening for therapeutic agents useful in the treatment of a disease comprising in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
 - i) contacting a test compound with a GSK3B polypeptide,
 - ii) detect binding of said test compound to said GSK3B polypeptide.
2. A method of screening for therapeutic agents useful in the treatment of a disease comprising in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
 - i) determining the activity of a GSK3B polypeptide at a certain concentration of a test compound or in the absence of said test compound,
 - ii) determining the activity of said polypeptide at a different concentration of said test compound.
3. A method of screening for therapeutic agents useful in the treatment of a disease comprising in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
 - i) determining the activity of a GSK3B polypeptide at a certain concentration of a test compound,
 - ii) determining the activity of a GSK3B polypeptide in the presence of a compound known to be a regulator of a GSK3B polypeptide.
4. The method of any of claims 1 to 3, wherein the step of contacting is in or at the surface of a cell.
5. The method of any of claims 1 to 3, wherein the cell is in vitro.
6. The method of any of claims 1 to 3, wherein the step of contacting is in a cell-free system.

7. The method of any of claims 1 to 3, wherein the polypeptide is coupled to a detectable label.
8. The method of any of claims 1 to 3, wherein the compound is coupled to a detectable label.
9. The method of any of claims 1 to 3, wherein the test compound displaces a ligand which is first bound to the polypeptide.
10. The method of any of claims 1 to 3, wherein the polypeptide is attached to a solid support.
11. The method of any of claims 1 to 3, wherein the compound is attached to a solid support.
12. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
 - i) contacting a test compound with a GSK3B polynucleotide,
 - ii) detect binding of said test compound to said GSK3B polynucleotide.
13. The method of claim 12 wherein the nucleic acid molecule is RNA.
14. The method of claim 12 wherein the contacting step is in or at the surface of a cell.
15. The method of claim 12 wherein the contacting step is in a cell-free system.
16. The method of claim 12 wherein polynucleotide is coupled to a detectable label.
17. The method of claim 12 wherein the test compound is coupled to a detectable label.
18. A method of diagnosing a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
 - i) determining the amount of a GSK3B polynucleotide in a sample taken from said mammal,
 - ii) determining the amount of GSK3B polynucleotide in healthy and/or diseased mammals.

19. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which binds to a GSK3B polypeptide.
- 5 20. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a GSK3B polypeptide.
- 10 21. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a GSK3B polypeptide, wherein said therapeutic agent is
- 15 i) a small molecule,
- ii) an RNA molecule,
- iii) an antisense oligonucleotide,
- iv) a polypeptide,
- v) an antibody, or
- 20 vi) a ribozyme.
22. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a GSK3B polynucleotide.
- 25 23. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a GSK3B polypeptide.

24. Use of regulators of a GSK3B for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal.
- 5 25. Method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
- 10 i) identifying a regulator of GSK3B,
- ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases in a mammal; and
- iii) combining of said regulator with an acceptable pharmaceutical carrier.
- 15 26. Use of a regulator of GSK3B for the regulation of GSK3B activity in a mammal having a disease comprised in a group of diseases consisting of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases.

Diagnostics and Therapeutics for Diseases Associated with Glycogen Synthase Kinase 3
Beta (GSK3B)

EPO - Munich
37

26. Feb. 2004

Abstract of Disclosure

The invention provides a human GSK3B which is associated with cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases. The invention also provides assays for the identification of compounds useful in the treatment or prevention of cardiovascular diseases, cancer, metabolic diseases, hematological diseases, inflammation, respiratory diseases, neurological diseases and urological diseases. The invention also features compounds which bind to and/or activate or inhibit the activity of GSK3B as well as pharmaceutical compositions comprising such compounds.

26. Feb. 2004

Fig. 1**SEQ ID NO: 1**

ATCATCTATATGTTAAATATCCGTGCCGATCTGTCTTGAAGGAGAAAT
ATATCGCTTGTTTTGTTTTTTATAGTATACAAAAGGAGTGAAAAGCCA
AGAGGACGAAGTCTTTTTCTTTTTCTTCTGTGGGAGAACTTAATGCTG
CATTTATCGTTAACCTAACACCCCAACATAAAGACAAAAGGAAGAAAA
GGAGGAAGGAAGGAAAAGGTGATTCGCGAAGAGAGTGATCATGTCAGG
GCGGCCCAGAACCACCTCCTTTGCGGAGAGCTGCAAGCCGGTGACAGCA
GCCTTCAGCTTTTGGCAGCATGAAAGTTAGCAGAGACAAGGACGGCAG
CAAGGTGACAACAGTGGTGGCAACTCCTGGGCAGGGTCCAGACAGGCC
ACAAGAAGTCAGCTATACAGACACTAAAGTGATTGGAAATGGATCATT
TGGTGTGGTATATCAAGCCAACTTTGTGATTCAGGAGAACTGGTCGC
CATCAAGAAAGTATTGCAGGACAAGAGATTTAAGAATCGAGAGCTCCA
GATCATGAGAAAGCTAGATCACTGTAACATAGTCCGATTGCGTTATTT
CTTCTACTCCAGTGGTGAGAAGAAAGATGAGGTCTATCTTAATCTGGT
GCTGGACTATGTTCCGGAAACAGTATACAGAGTTGCCAGACACTATAG
TCGAGCCAAACAGACGCTCCCTGTGATTTATGTCAAGTTGTATATGTA
TCAGCTGTTCCGAAGTTTAGCCTATATCCATTCCTTTGGAAATCTGCCA
TCGGGATATTAAACCGCAGAACCTCTTGTTGGATCCTGATACTGCTGT
ATTAAAACTCTGTGACTTTGGAAGTGCAAAGCAGCTGGTCCGAGGAGA
ACCCAATGTTTTCGTATATCTGTTCTCGGTACTATAGGGCACCAGAGTT
GATCTTTGGAGCCACTGATTATACCTCTAGTATAGATGTATGGTCTGC
TGGCTGTGTGTTGGCTGAGCTGTTACTAGGACAACCAATATTTCCAGG
GGATAGTGGTGTGGATCAGTTGGTAGAAATAATCAAGGTCCTGGGAAC
TCCAACAAGGGAGCAAATCAGAGAAATGAACCCAACTACACAGAATT
TAAATTCCCTCAAATTAAGGCACATCCTTGGACTAAGGATTCGTCAGG
AACAGGACATTTACCTCAGGAGTGCGGGTCTTCCGACCCCGAACTCC
ACCGGAGGCAATTGCACTGTGTAGCCGTCTGCTGGAGTATACACCAAC
TGCCCGACTAACACCACTGGAAGCTTGTGCACATTCATTTTTTTGATGA
ATTACGGGACCCAAATGTCAAACCTACCAAATGGGCGAGACACACCTGC

ACTCTTCAACTTCACCACTCAAGAACTGTCAAGTAATCCACCTCTGGC
TACCATCCTTATTCCTCCTCATGCTCGGATTCAAGCAGCTGCTTCAAC
CCCCACAAATGCCACAGCAGCGTCAGATGCTAATACTGGAGACCGTGG
ACAGACCAATAATGCTGCTTCTGCATCAGCTTCCAACCTCCACCTGAAC
AGTCCCGAGCAGCCAGCTGCACAGGAAAAACCACCAGTTACTTGAGTG
TCACTCAGCAACACTGGTCACGTTTGGAAAGAATATTAAAAAAAAAAAA
AAAAAAA

Fig. 2

SEQ ID NO: 2

MSGRPRTTSFAESCKPVQQPSAFGSMKVS RDKDGSKVTTVVATPGQGP
DRPQEVSYTDTKVIGNGSFGVVYQAKLCDSGELVAIKKVLQDKRFKNR
ELQIMRKLDHCNIVRLRYFFYS SGEKKDEVYLNLVLDYVPETVYRVAR
HYSRAKQTL PVIYVKLYMYQLFRSLAYIHSFGICH RDIKPQNLLDPD
TAVLKLCDFGSAKQLVRGEPNVSYICSRY YRAPELIFGATDYTSSIDV
WSAGCVLAELLLGQPIFP GDSGVDQLVEIIKVLGTP TREQIREMNP NY
TEFKFPQIKAHPWTKDSSGTGHFTSGVRVFRP RTPPEAIALCSR LLEY
TPTARLTPLEACAHSFFDEL RDPNVKLPNGRDT PALFNFTTQELSSNP
PLATILIPPHARIQAAASTPTNATAASDANTGDRGQTNNAASASASNS
T

Fig. 3

SEQ ID NO: 3

5' -TTCCAGGGGATAGTGGTGTG-3'

Fig. 4

SEQ ID NO: 4

5' -TTTGCTCCCTTGTTGGAGTT-3'

Fig. 5

SEQ ID NO: 5

5' -TCAGTTGGTAGAAATAATCAAGGTCCTGGG-3'

26. Feb. 2004

SEQUENCE LISTING

<110> Bayer HealthCare AG

<120> Diagnostics and Therapeutics for Diseases Associated with
Glycogen Synthase Kinase 3 beta (GSK3B)

<130> BHC 04 01 040

<160> 5

<170> PatentIn version 3.2

<210> 1

<211> 1639

<212> DNA

<213> Homo sapiens

<400> 1

```
atcatctata tgttaaataat ccgtgccgat ctgtcttgaa ggagaaatat atcgcttgtt      60
ttgtttttta tagtatacaa aaggagtga aagccaagag gacgaagtct ttttcttttt      120
cttctgtggg agaacttaat gctgcattta tcgttaacct aacaccccaa cataaagaca      180
aaaggaagaa aaggaggaag gaaggaaaag gtgattcgcg aagagagtga tcatgtcagg      240
gcggcccaga accacctcct ttgcggagag ctgcaagccg gtgcagcagc cttcagcttt      300
tggcagcatg aaagttagca gagacaagga cggcagcaag gtgacaacag tggaggcaac      360
tcctgggcag ggtccagaca ggccacaaga agtcagctat acagacacta aagtgattgg      420
aatggatca tttggtgtgg tatatcaagc caaactttgt gattcaggag aactggtcgc      480
catcaagaaa gtattgcagg acaagagatt taagaatcga gagctccaga tcatgagaaa      540
gctagatcac tgtaacatag tccgattgcg ttattttctt tactccagtg gtgagaagaa      600
agatgagggtc tatcttaatc tgggtgctgga ctatgttccg gaaacagtat acagagttgc      660
cagacactat agtcgagcca aacagacgct ccctgtgatt tatgtcaagt tgtatatgta      720
tcagctgttc cgaagttag cctatatcca ttcttttggg atctgccatc gggatattaa      780
accgcagaac ctcttgttgg atcctgatac tgctgtatta aaactctgtg actttggaag      840
tgcaaagcag ctggtccgag gagaacccaa tgtttcgtat atctgttctc ggtactatag      900
ggcaccagag ttgatctttg gagccactga ttatacctct agtatagatg tatggtctgc      960
tggctgtgtg ttggctgagc tgttactagg acaaccaata tttccagggg atagtgggtg      1020
ggatcagttg gtagaaataa tcaaggctct gggaaactcca acaagggagc aaatcagaga      1080
aatgaaccca aactacacag aatttaaatt ccctcaaatt aaggcacatc cttggactaa      1140
ggattcgtca ggaacaggac atttcacctc aggagtgcgg gtcttccgac cccgaactcc      1200
accggaggca attgcactgt gtagccgtct gctggagtat acaccaactg cccgactaac      1260
accactggaa gcttgtgcac attcattttt tgatgaatta cgggacccaa atgtcaaact      1320
accaaattgg cgagacacac ctgcactott caacttcacc actcaagaac tgtcaagtaa      1380
tccacctctg gctaccatcc ttattcctcc tcatgctcgg attcaagcag ctgcttcaac      1440
ccccacaaat gccacagcag cgtcagatgc taatactgga gaccgtggac agaccaataa      1500
tgctgcttct gcacagctt ccaactccac ctgaacagtc ccgagcagcc agctgcacag      1560
```

gaaaaaccac cagttacttg agtgtcactc agcaacactg gtcacgtttg gaaagaatat 1620
 taaaaaaaaa aaaaaaaaaa 1639

<210> 2

<211> 433

<212> PRT

<213> Homo sapiens

<400> 2

Met Ser Gly Arg Pro Arg Thr Thr Ser Phe Ala Glu Ser Cys Lys Pro
 1 5 10 15
 Val Gln Gln Pro Ser Ala Phe Gly Ser Met Lys Val Ser Arg Asp Lys
 20 25 30
 Asp Gly Ser Lys Val Thr Thr Val Val Ala Thr Pro Gly Gln Gly Pro
 35 40 45
 Asp Arg Pro Gln Glu Val Ser Tyr Thr Asp Thr Lys Val Ile Gly Asn
 50 55 60
 Gly Ser Phe Gly Val Val Tyr Gln Ala Lys Leu Cys Asp Ser Gly Glu
 65 70 75 80
 Leu Val Ala Ile Lys Lys Val Leu Gln Asp Lys Arg Phe Lys Asn Arg
 85 90 95
 Glu Leu Gln Ile Met Arg Lys Leu Asp His Cys Asn Ile Val Arg Leu
 100 105 110
 Arg Tyr Phe Phe Tyr Ser Ser Gly Glu Lys Lys Asp Glu Val Tyr Leu
 115 120 125
 Asn Leu Val Leu Asp Tyr Val Pro Glu Thr Val Tyr Arg Val Ala Arg
 130 135 140
 His Tyr Ser Arg Ala Lys Gln Thr Leu Pro Val Ile Tyr Val Lys Leu
 145 150 155 160
 Tyr Met Tyr Gln Leu Phe Arg Ser Leu Ala Tyr Ile His Ser Phe Gly
 165 170 175
 Ile Cys His Arg Asp Ile Lys Pro Gln Asn Leu Leu Leu Asp Pro Asp
 180 185 190
 Thr Ala Val Leu Lys Leu Cys Asp Phe Gly Ser Ala Lys Gln Leu Val
 195 200 205
 Arg Gly Glu Pro Asn Val Ser Tyr Ile Cys Ser Arg Tyr Tyr Arg Ala
 210 215 220
 Pro Glu Leu Ile Phe Gly Ala Thr Asp Tyr Thr Ser Ser Ile Asp Val
 225 230 235 240
 Trp Ser Ala Gly Cys Val Leu Ala Glu Leu Leu Leu Gly Gln Pro Ile
 245 250 255
 Phe Pro Gly Asp Ser Gly Val Asp Gln Leu Val Glu Ile Ile Lys Val
 260 265 270
 Leu Gly Thr Pro Thr Arg Glu Gln Ile Arg Glu Met Asn Pro Asn Tyr
 275 280 285

- 3 -

Thr Glu Phe Lys Phe Pro Gln Ile Lys Ala His Pro Trp Thr Lys Asp
 290 295 300
 Ser Ser Gly Thr Gly His Phe Thr Ser Gly Val Arg Val Phe Arg Pro
 305 310 315 320
 Arg Thr Pro Pro Glu Ala Ile Ala Leu Cys Ser Arg Leu Leu Glu Tyr
 325 330 335
 Thr Pro Thr Ala Arg Leu Thr Pro Leu Glu Ala Cys Ala His Ser Phe
 340 345 350
 Phe Asp Glu Leu Arg Asp Pro Asn Val Lys Leu Pro Asn Gly Arg Asp
 355 360 365
 Thr Pro Ala Leu Phe Asn Phe Thr Thr Gln Glu Leu Ser Ser Asn Pro
 370 375 380
 Pro Leu Ala Thr Ile Leu Ile Pro Pro His Ala Arg Ile Gln Ala Ala
 385 390 395 400
 Ala Ser Thr Pro Thr Asn Ala Thr Ala Ala Ser Asp Ala Asn Thr Gly
 405 410 415
 Asp Arg Gly Gln Thr Asn Asn Ala Ala Ser Ala Ser Ala Ser Asn Ser
 420 425 430
 Thr

<210> 3
 <211> 20
 <212> DNA
 <213> artificial sequence

<220>
 <223> forward primer

<400> 3
 ttccagggga tagtggtgtg

20

<210> 4
 <211> 20
 <212> DNA
 <213> artificial sequence

<220>
 <223> reverse primer

<400> 4
 tttgctccct tgttggagtt

20

<210> 5
 <211> 30

<212> DNA

<213> artificial sequence

<220>

<223> probe

<400> 5

tcagttggta gaaataatca aggtcctggg